

Characterization of bacterial cellulose-neem-hypericum oil wound care paste *in vitro* and in *Galleria mellonella* *in vivo* model

Stefania Villani¹, Sanosh Kunjalukkal Padmanabahn¹, Mariangela Stoppa², Rossella Nisi², Matteo Calcagnile³, Pietro Alifano³, Christian Demitri³, and Antonio Licciulli¹

¹Department of Engineering for Innovation (DII) – Università del Salento, Lecce, Italy

²Biofaber Srl – Brindisi, Italy

³Department of Experimental Medicine – Università del Salento, Lecce, Italy

Surgical wound infections represent a global health emergency because of the spread of antibiotic-resistant bacteria. In this context, nanotechnologies help the optimization of strategies for managing skin lesions with antibacterial and antibiofilm activity.

In this work, we developed a biobased formulation made with bacterial cellulose nanofibers obtained through a green fermentation process, combined with a mixture of neem and hypericum oils, commonly used in the cosmetic field (1).

The resulting bacterial cellulose-based paste has been physicochemical characterized. *In vitro* antibacterial assays showed that the bacterial cellulose-based paste has biofilm detachment capabilities against *Pseudomonas aeruginosa* and *Staphylococcus aureus*. The efficacy of the treatment was also evaluated *in vivo* using *Galleria mellonella* larvae as a burn wound infection model, observing an increase in the survival percentage of the injured and infected larvae treated with our formulation, compared to untreated individuals. The innovative aspect of this formulation is the detachment action of bacterial biofilms, which contributes to a faster tissue regeneration process without inducing antibiotic resistance. Furthermore, in this study, the involvement of *G. mellonella* larvae as a model of burn wound infection allowed a preliminary evaluation of the safety and efficacy of the formulation, contributing to the development of a more complex system than traditional *in vitro* ones, but limiting the involvement of mammalian models, in line with the 3Rs principle.

These results, supported by further investigation, could lead to the development of a biobased formulation to be applied on skin lesions for antibacterial treatments.

(1) <https://doi.org/10.1016/j.carpta.2024.100431>