

Fine gradings on the simple Lie algebras of type E

Cristina Draperⁱ

Department of Applied Mathematics, University of Málaga
cdf@uma.es

Alberto Elduqueⁱⁱ

Department of Mathematics, University of Zaragoza
elduque@unizar.es

Abstract. Some fine gradings on the exceptional Lie algebras \mathfrak{e}_6 , \mathfrak{e}_7 and \mathfrak{e}_8 are described. This list tries to be exhaustive.

Keywords: Grading; simple; exceptional; Lie algebra; type E .

MSC 2010 classification: primary 17B70, secondary 17B25

Introduction

Gradings by groups have played a key role in the study of Lie algebras, and contributed to understand their structural properties. Several families of examples can be found in [23]. To begin with, the root space decomposition of a complex semisimple Lie algebra is a grading by the group \mathbb{Z}^r , with r the rank of the Lie algebra. Any grading by a torsion-free abelian group is equivalent to a coarsening of such root space decomposition, and these gradings have been extensively used in representation theory. Gradings by not necessarily reduced root systems are very nice examples of this situation. In particular, gradings by the integers have had frequent applications to physics, and they are specially relevant in algebraic contexts: if J is a Jordan algebra, the Tits-Kantor-Koecher construction applied to J is a \mathbb{Z} -graded Lie algebra $L = L_{-1} \oplus L_0 \oplus L_1$ with $L_1 = J$, and the product in J can be recovered from the one in L . Some other Jordan systems are related to ‘longer’ \mathbb{Z} -gradings too.

Gradings by groups with torsion are also ubiquitous: gradings by cyclic groups and the corresponding finite order automorphisms are described by Kac

ⁱSupported by the Spanish Ministerio de Economía y Competitividad—Fondo Europeo de Desarrollo Regional (FEDER) MTM 2010-15223 and by the Junta de Andalucía grants FQM-336, FQM-1215 and FQM-246.

ⁱⁱSupported by the Spanish Ministerio de Economía y Competitividad—Fondo Europeo de Desarrollo Regional (FEDER) MTM2010-18370-C04-02 and by the Diputación General de Aragón—Fondo Social Europeo (Grupo de Investigación de Álgebra).

<http://siba-ese.unisalento.it/> © 2014 Università del Salento

[38]. They are intimately related to (infinite-dimensional) Kac-Moody Lie algebras and gradings by \mathbb{Z} on them. In Differential Geometry, they are connected with symmetric spaces and their generalizations. Gradings by finite abelian groups are related to Lie color algebras (a generalization of Lie superalgebras), and sometimes to Lie algebra contractions. An interested reader can consult [23, Introduction] for these and other examples, as well as for applications and references. Also a good and deep compilation of results can be found in [46, Chapter 3, §3], which deals with the relationship between gradings on complex semisimple Lie algebras and automorphisms, and exhibit a wide variety of examples of gradings.

J. Patera and H. Zassenhaus, convinced about the relevance of gradings, initiated in [47] a systematic study of gradings on Lie algebras, emphasizing the role of the so called fine gradings (gradings which cannot be further refined). Since then, a considerable number of authors have been trying to obtain a classification of gradings on the simple Lie algebras (see, e.g. [33, 9, 21, 15, 16, 17]), which has culminated in the recent monograph [23], where gradings on the classical simple Lie algebras and on the exceptional simple Lie algebras of types G_2 and F_4 are thoroughly studied. However, there is still work to be done. On one hand, not much is known about gradings on solvable or nilpotent Lie algebras. On the other hand, the classification of gradings (for instance, the classification of fine gradings up to equivalence) is not yet finished for the complex exceptional simple Lie algebras of types E_7 and E_8 (denoted by \mathfrak{e}_7 and \mathfrak{e}_8), and this is also the case for the simple Lie algebras of types D_4 and E_r ($r = 6, 7, 8$) over algebraically closed fields of prime characteristic. Over the real numbers, even the classification of fine gradings for the classical Lie algebras is missing, although many low-dimensional cases have been considered.

Our goal is the classification of the fine gradings on the exceptional Lie algebras \mathfrak{e}_7 and \mathfrak{e}_8 over an algebraically closed field \mathbb{F} of characteristic zero, and hence to finish the classification of fine gradings on simple Lie algebras over the complex numbers. (Note the result in [17, Proposition 2], which shows that the complex case yields a solution over arbitrary algebraically closed fields of characteristic 0.)

This goal has not been reached yet. The purpose of this paper is to describe a list of known fine gradings, which are compiled in our Main Theorem (Theorem 3.3) on the exceptional simple Lie algebras of type E . Most of these gradings make sense in much more general contexts but, to avoid confusion, we will restrict ourselves to an algebraically closed ground field \mathbb{F} of characteristic zero. The list exhausts the fine gradings, up to equivalence, on \mathfrak{e}_6 , and we conjecture that it also exhausts them for \mathfrak{e}_7 and \mathfrak{e}_8 . This has been announced in [23, Figure 6.2], although not all the descriptions there coincide with ours. Further

details will be provided here too.

In our setting, a fine grading is the eigenspace decomposition relative to a maximal quasitorus (that is, a maximal abelian diagonalizable, or MAD, subgroup) of the group of automorphisms. This means that fine gradings are related to ‘large’ abelian groups of symmetries. Hence our goal is equivalent to the classification of the MAD-subgroups, up to conjugation, of $\text{Aut}(\mathfrak{e}_7)$ and $\text{Aut}(\mathfrak{e}_8)$.

Some of these MAD-subgroups have appeared in the literature. For instance, Griess describes in [31] the maximal elementary p -subgroups of the groups E_6 , E_7 and E_8 . A larger class of abelian subgroups (not just MAD-subgroups) is studied in [52]. The \mathbb{Z}_6^3 -subgroup of E_8 has been considered by Hang and Vogan in [32, pp. 22-25]. Also, Alekseevskii described the Jordan finite commutative subgroups of the groups E_6 , E_7 and E_8 in [1]. These include a MAD-subgroup of E_8 isomorphic to \mathbb{Z}_5^3 . This subgroup has gained some attention lately, as it appeared in a talk by Kostant (see [42]) about the controversial paper by the physicist Lisi [44], which proposed a theory to go beyond the Standard Model in that it unifies all 4 forces of nature by using as gauge group the exceptional Lie group E_8 . Kostant’s talk, strictly mathematical, dealt about an elaboration of the mathematics of E_8 in order to refute Lisi’s Theory. This is one more evidence of the fascination produced by the richness of E_8 , and shows the relevance of understanding as much as possible about this group and its tangent Lie algebra. Note that not even the finite abelian maximal groups are conveniently well known (consult the recent work [14]). In terms of gradings, some of our descriptions have appeared in [19], which uses gradings on composition algebras to construct some *nice* gradings on exceptional algebras (e.g. a \mathbb{Z}_3^5 -grading and a \mathbb{Z}_2^8 -grading on \mathfrak{e}_8).

This paper then gathers a lot of known material, and describes it in a homogeneous way. It is an expanded version of the talk presented by the first author in the conference *Advances in Group Theory and Applications 2013*.

The paper is structured as follows. First there is a section to recall the background: basic concepts about gradings and their connection with groups of automorphisms, and some algebraic structures involved in the description of the exceptional Lie algebras: composition (and symmetric composition) algebras, Jordan algebras and its generalizations, and structurable algebras. Second, we present several models of the exceptional Lie algebras and constructions leading to them. After reviewing slightly how the exceptional Lie algebras emerged, we focus mainly on three constructions due to Elduque, Kantor and Steinberg, as they provide a convenient way to describe the fine gradings we are interested in. Finally, the third section describes some fine gradings on \mathfrak{e}_6 , \mathfrak{e}_7 and \mathfrak{e}_8 starting from gradings on the ‘coordinate algebras’ involved in the constructions above. (These algebras are usually much simpler than \mathfrak{e}_6 , \mathfrak{e}_7 and \mathfrak{e}_8 .) Fourteen fine gradings will be given on each of these simple Lie algebras. The conjecture

arises immediately of whether this is the complete list of fine gradings, up to equivalence, for these algebras.

For the sake of clarity of the exposition, in what follows the ground field \mathbb{F} will be assumed to be algebraically closed of characteristic 0, even though many results are valid in more general contexts.

1 Preliminaries

1.1 Gradings and automorphisms

We begin by recalling the basics about gradings. Let \mathcal{A} be a finite-dimensional algebra (not necessarily associative) over \mathbb{F} , and let G be an abelian group.

Definition 1.1. A G -grading Γ on \mathcal{A} is a vector space decomposition

$$\Gamma : \mathcal{A} = \bigoplus_{g \in G} \mathcal{A}_g$$

such that $\mathcal{A}_g \mathcal{A}_h \subseteq \mathcal{A}_{g+h}$ for all $g, h \in G$.

Once such a decomposition is fixed, the algebra \mathcal{A} will be called a G -graded algebra, the subspace \mathcal{A}_g will be referred to as the *homogeneous component of degree g* and its nonzero elements will be called the *homogeneous elements of degree g* . The *support* is the set $\text{supp } \Gamma := \{g \in G \mid \mathcal{A}_g \neq 0\}$.

Definition 1.2. If $\Gamma : \mathcal{A} = \bigoplus_{g \in G} \mathcal{A}_g$ and $\Gamma' : \mathcal{A} = \bigoplus_{h \in H} \mathcal{A}_h$ are gradings by two abelian groups G and H , Γ is said to be a *refinement* of Γ' (or Γ' a *coarsening* of Γ) if for any $g \in G$ there is $h \in H$ such that $\mathcal{A}_g \subseteq \mathcal{A}_h$. In other words, any homogeneous component of Γ' is the direct sum of some homogeneous components of Γ . A refinement is *proper* if some inclusion $\mathcal{A}_g \subseteq \mathcal{A}_h$ is proper. A grading is said to be *fine* if it admits no proper refinement.

Definition 1.3. Let Γ be a G -grading on \mathcal{A} and Γ' an H -grading on another algebra \mathcal{B} , with supports, respectively, S and T . Then Γ and Γ' are said to be *equivalent* if there is an algebra isomorphism $\varphi : \mathcal{A} \rightarrow \mathcal{B}$ and a bijection $\alpha : S \rightarrow T$ such that $\varphi(\mathcal{A}_s) = \mathcal{B}_{\alpha(s)}$ for all $s \in S$. Any such φ is called an *equivalence* of Γ and Γ' .

The study of gradings is based on classifying fine gradings up to equivalence, because any grading is obtained as a coarsening of some fine one. We will make use of the following invariant by equivalences:

Definition 1.4. The *type* of a grading Γ is the sequence of numbers (h_1, \dots, h_r) where h_i is the number of homogeneous components of dimension i , with $i = 1, \dots, r$ and $h_r \neq 0$. Obviously, $\dim \mathcal{A} = \sum_{i=1}^r i h_i$.

Given a group grading Γ on an algebra \mathcal{A} , there are many groups G such that Γ , regarded as a decomposition into a direct sum of subspaces such that the product of any two of them lies in a third one, can be realized as a G -grading, but there is one distinguished group among them ([47]). Define $U(\Gamma)$ as the abelian group generated by $S = \text{supp } \Gamma$ with defining relations $s_1 s_2 = s_3$ whenever $0 \neq \mathcal{A}_{s_1} \mathcal{A}_{s_2} \subset \mathcal{A}_{s_3}$ ($s_i \in S$). It is called the *universal group of Γ* , since it verifies that, for any other realization of Γ as a G -grading, there exists a unique homomorphism $U(\Gamma) \rightarrow G$ that restricts to identity on $\text{supp } \Gamma$. All the gradings throughout this work will be considered by their universal groups.

The classification of fine gradings on \mathcal{A} , up to equivalence, is the same as the classification of maximal diagonalizable subgroups (i.e., maximal quasitori) of $\text{Aut}(\mathcal{A})$, up to conjugation (see e.g. [47]). More precisely, given a G -grading on the algebra $\mathcal{A} = \bigoplus_{g \in G} \mathcal{A}_g$, any χ belonging to the group of characters $\hat{G} = \text{Hom}(G, \mathbb{F}^\times)$, acts as an automorphism of \mathcal{A} by means of $\chi.x = \chi(g)x$ for any $g \in G$ and $x \in \mathcal{A}_g$. In case G is the universal group of the grading, this allows us to identify \hat{G} with a quasitorus (the direct product of a torus and a finite subgroup) of the algebraic group $\text{Aut}(\mathcal{A})$. This quasitorus is the subgroup $\text{Diag}(\Gamma)$ consisting of the automorphisms φ of \mathcal{A} such that the restriction of φ to any homogeneous component is the multiplication by a (nonzero) scalar. (See [46, Chapter 3, §3] or [23, §1.4].) Conversely, given a quasitorus Q of $\text{Aut}(\mathcal{A})$, then Q induces a \hat{Q} -grading on \mathcal{A} , where $\mathcal{A}_g = \{x \in \mathcal{A} \mid \chi(x) = g(\chi)x \forall \chi \in Q\}$ for any $g \in \hat{Q}$. In this way the fine gradings on \mathcal{A} , up to equivalence, correspond to the conjugacy classes in $\text{Aut}(\mathcal{A})$ of the maximal abelian diagonalizable subgroups of $\text{Aut}(\mathcal{A})$.

1.2 Related structures

We will recall here some algebraic structures involved in the constructions of the exceptional Lie algebras. A very nice introduction to nonassociative algebras can be found in [48], but the necessary material of composition algebras and Jordan algebras is included here for completeness, as well as material about symmetric composition algebras and structurable algebras, which are not so well known.

1.2.1 Composition algebras

A Hurwitz algebra over \mathbb{F} is a unital algebra C endowed with a nonsingular quadratic form $q: C \rightarrow \mathbb{F}$ which is multiplicative, that is, $q(xy) = q(x)q(y)$. This form q is usually called the *norm*. Each element $a \in C$ satisfies

$$a^2 - t_C(a)a + q(a)1 = 0,$$

where $t_C(a) = q(a+1) - q(a) - 1$ is called the *trace*. Denote $C_0 = \{a \in C \mid t_C(a) = 0\}$ the subspace of traceless elements. Note that $[a, b] = ab - ba \in C_0$ for any $a, b \in C$, since $t_C(ab) = t_C(ba)$. The map $- : C \rightarrow C$ given by $\bar{a} = t_C(a)1 - a$ is an involution and $q(a) = a\bar{a}$ holds.

There are Hurwitz algebras only in dimensions 1, 2, 4 and 8 (see e.g. [53]). Moreover, under our hypothesis on the field, there is only one (up to isomorphism) Hurwitz algebra of each possible dimension, namely:

- the ground field \mathbb{F} , with $q(a) = a^2$;
- $\mathbb{F} \times \mathbb{F}$, with componentwise product and norm given by $q(a, b) = ab$;
- $\text{Mat}_{2 \times 2}(\mathbb{F})$, with the usual matrix product and norm given by the determinant;
- the split Cayley algebra over \mathbb{F} . This algebra can be characterized by the existence of a basis $\{e_1, e_2, u_1, u_2, u_3, v_1, v_2, v_3\}$, which we call *standard basis*, with multiplication given by

$$\begin{aligned} e_1 u_j &= u_j = u_j e_2, & u_i v_i &= e_1, & u_i u_{i+1} &= v_{i+2} = -u_{i+1} u_i, \\ e_2 v_j &= v_j = v_j e_1, & v_i u_i &= e_2, & -v_i v_{i+1} &= u_{i+2} = v_{i+1} v_i, \end{aligned}$$

all the remaining products being 0, and the polar form of the norm (also denoted by q) of two basic elements is zero except for $q(e_1, e_2) = 1 = q(u_i, v_i)$, $i = 1, 2, 3$.

With the exception of the Cayley algebra, all of these are associative. The Cayley algebra is not associative but alternative (the algebra generated by any pair of elements is associative). We will use the notations \mathbb{F} , \mathcal{K} , \mathcal{Q} (usually called *quaternion algebra*) and \mathcal{C} (usually called *octonion algebra*) for each of these algebras, respectively.

Recall that for any $a, b \in C$, the endomorphism

$$d_{a,b} = [l_a, l_b] + [l_a, r_b] + [r_a, r_b]$$

is a derivation of C for $l_a(b) = ab$ and $r_a(b) = ba$. This will be instrumental to construct Lie algebras from Hurwitz algebras.

1.2.2 Jordan algebras

A *Jordan algebra* is a commutative (nonassociative) algebra satisfying the Jordan identity

$$(x^2 y)x = x^2(yx).$$

This kind of algebras were introduced by Jordan in 1933 to formalize the notion of an algebra of observables in quantum mechanics. Such line of research was abandoned time ago, but Jordan algebras have found a range of applications because of their relationship to Lie algebras. A standard reference is [36].

If A is an associative algebra (with multiplication denoted by juxtaposition) and we consider the new product on A given by

$$x \circ y = \frac{1}{2}(xy + yx),$$

then (A, \circ) is a Jordan algebra, denoted by A^+ . A Jordan algebra which is a subalgebra of A^+ for some associative algebra A , is called a *special* Jordan algebra, and otherwise it is called *exceptional*. If $(A, -)$ is an associative algebra with involution, then the set of hermitian elements $H(A, -) = \{a \in A \mid \bar{a} = a\}$ is a subalgebra of A^+ (not of A), and hence it is a special Jordan algebra.

In particular, if C is an associative Hurwitz algebra with involution given by $-$, the algebra $H_n(C, *) = \{x = (x_{ij}) \in \text{Mat}_{n \times n}(C) \mid x_{ij} = \bar{x}_{ji}\}$ is a Jordan algebra for any $n \geq 3$. (For $n = 1$ or $n = 2$ this is also true, but in a trivial way, so we will assume $n \geq 3$.) It is proved in [36] that if C is the Cayley algebra, $H_n(C, *)$ is a Jordan algebra if and only if $n = 3$. Besides, this is the only exceptional Jordan algebra, which is called the Albert algebra, and will be denoted by \mathbb{A} .

If $J = H_n(C, *)$ for some Hurwitz algebra and some n , consider the linear map $t_J: J \rightarrow \mathbb{F}$ given by $t_J(x) = \frac{\text{tr}(x)}{n} = \frac{\sum_{i=1}^n x_{ii}}{n}$. This map is called the *normalized trace* and it is the only linear map such that $t_J(1) = 1$ and $t_J((xy)z) = t_J(x(yz))$ for any $x, y, z \in J$. Thus we have a decomposition $J = \mathbb{F}1 \oplus J_0$, for $J_0 = \{x \in J \mid t_J(x) = 0\}$, since $x * y = xy - t_J(xy)1 \in J_0$. In particular we have a commutative multiplication $*$ defined in J_0 .

If J is a Jordan algebra and $R_x: J \rightarrow J, y \mapsto yx$ is the multiplication operator, observe that

$$[[R_x, R_y], R_z] = R_{(yz)x - y(zx)} \tag{1}$$

for any $x, y, z \in J$, and thus, the *structure algebra* $\mathfrak{Str}(J)$, or Lie algebra generated by the multiplication operators, coincides with $R_J + [R_J, R_J]$ (this sum is direct if J is unital). It is also a consequence of Equation (1) that $[R_J, R_J]$ is an ideal of the Lie algebra of derivations $\text{Der}(J) = \{d \in \mathfrak{gl}(J) \mid d(xy) = d(x)y + xd(y) \forall x, y \in J\}$. The algebra generated by the traceless multiplication operators $\{R_x \mid x \in J_0\}$ is called the *inner structure algebra* and it also coincides with $R_{J_0} + [R_J, R_J]$.

In case $J = \mathbb{A}$ is the Albert algebra, $\text{Der}(\mathbb{A})$ is simple [37, Theorem 3], so in particular every derivation is inner ($\text{Der}(\mathbb{A}) = [R_{\mathbb{A}}, R_{\mathbb{A}}]$). The inner structure

algebra is simple too [37, Theorem 4], and these provide our first two models of \mathfrak{f}_4 and \mathfrak{e}_6 . Moreover, any element $x \in \mathbb{A}$ satisfies a cubic equation

$$x^3 - T(x)x^2 + Q(x)x - N(x)1 = 0, \quad (2)$$

for the scalars $T(x) = \text{tr}(x)$, $Q(x) = \frac{1}{2}((T(x))^2 - T(x^2))$ and $N(x) = \frac{1}{6}((T(x))^3 - 3T(x)T(x^2) + 2T(x^3))$. The cubic form N is also closely related to \mathfrak{e}_6 .

1.2.3 Symmetric composition algebras

A *symmetric composition algebra* is a triple $(S, *, q)$, where $(S, *)$ is a (nonassociative) algebra over \mathbb{F} with multiplication denoted by $x * y$ for $x, y \in S$, and where $q: S \rightarrow \mathbb{F}$ is a regular quadratic form verifying

$$\begin{aligned} q(x * y) &= q(x)q(y), \\ q(x * y, z) &= q(x, y * z), \end{aligned}$$

for any $x, y, z \in S$, where $q(x, y) = q(x + y) - q(x) - q(y)$ is the polar form of q .

Example 1.5. Let C be a Hurwitz algebra with norm q . The same vector space with new product

$$x * y = \bar{x}\bar{y}$$

for any $x, y \in C$ is a symmetric composition algebra for the same norm, called the *para-Hurwitz algebra* attached to the Hurwitz algebra C . We will denote it by $pC = (C, *, q)$. Note that the unit of C becomes a *paraunit* in pC , that is, an element e such that $e * x = x * e = q(e, x)e - x$.

Example 1.6. The *Okubo algebra*, or pseudo-octonion algebra, is the algebra defined on the subspace of trace 0 matrices of degree 3: $\text{Ok} = (\text{Mat}_{3 \times 3}(\mathbb{F})_0, *, q)$ with multiplication

$$x * y = \omega xy - \omega^2 yx - \frac{\omega - \omega^2}{3} \text{tr}(xy)1 \quad (3)$$

and norm $q(x) := \frac{1}{6} \text{tr}(x^2)$, for $x, y \in \text{Mat}_{3 \times 3}(\mathbb{F})_0$, where ω is a primitive cubic root of 1. This algebra is a symmetric composition algebra, but it does not have an identity element (and it is not alternative). It was introduced by Okubo in [45], who was working in Particle Physics and the symmetry given by the compact group $\text{SU}(3)$ (the real algebra $\mathfrak{su}(3) = \{x \in \mathfrak{sl}(3, \mathbb{C}) \mid x^* = -x\}$, for $*$ the unitary involution, is closed for the product given in Equation (3)).

The classification of the symmetric composition algebras was obtained in [24].

Theorem 1.7. *Every symmetric composition algebra over \mathbb{F} (algebraically closed) is isomorphic either to a para-Hurwitz algebra or to the Okubo algebra. That is, there are only five symmetric composition algebras (up to isomorphism), namely, $p\mathbb{F}$, $p\mathcal{K}$, $p\mathcal{Q}$, $p\mathcal{C}$ and Ok.*

1.2.4 Structurable algebras

Let $(A, -)$ be a unital algebra with involution “ $-$ ”. Denote the multiplication in A by juxtaposition. For $x, y \in A$, consider the linear operator $V_{x,y} \in \text{End}(A)$ given by $V_{x,y}(z) = (x\bar{y})z + (z\bar{y})x - (z\bar{x})y \equiv \{x, y, z\}$. The algebra A is called a *structurable algebra* in case the identity

$$\{x, y, \{z, w, v\}\} = \{\{x, y, z\}, w, v\} - \{z, \{y, x, w\}, v\} + \{z, w, \{x, y, v\}\}$$

is satisfied for any $x, y, z, w, v \in A$, or equivalently,

$$[V_{x,y}, V_{z,w}] = V_{V_{x,y}z, w} - V_{z, V_{y,x}w}.$$

The reader may consult [2] for the definition and properties of structurable algebras.

Example 1.8. Any (unital) associative algebra with involution $(A, -)$ is a structurable algebra.

The space $\mathfrak{Inst}(A, -) = \{\sum_i V_{x_i, y_i} \mid x_i, y_i \in A\}$ is a subalgebra of the Lie algebra $\mathfrak{gl}(A)$, called the *inner structure algebra* of $(A, -)$. The map

$$\varepsilon: \mathfrak{Inst}(A, -) \rightarrow \mathfrak{Inst}(A, -), \quad \varepsilon(V_{x,y}) = -V_{y,x}$$

if $x, y \in A$, is an involutive automorphism of this Lie algebra. Thus $\mathfrak{Inst}(A, -)$ turns out to be \mathbb{Z}_2 -graded. The elements in $H(A, -) = \{x \in A \mid \bar{x} = x\}$ and $S(A, -) = \{x \in A \mid \bar{x} = -x\}$ are called hermitian and skew-hermitian respectively. It follows that $\mathfrak{Inst}(A, -)_{\bar{0}} = (\mathfrak{Inst}(A, -) \cap \text{Der}(A, -)) \oplus V_{S(A, -), 1}$ and $\mathfrak{Inst}(A, -)_{\bar{1}} = V_{H(A, -), 1}$, where $\text{Der}(A, -)$ denotes the set of derivations of A that commute with the involution $-$.

Example 1.9. If J is a Jordan algebra, then $(J, -)$ is a structurable algebra with the involution $-$ given by the identity map. In this case $V_{x,y} = R_{xy} + [R_x, R_y]$ for $x, y \in J$, where R_x is the multiplication operator by x . In this sense, the inner structure algebra of $(J, -)$ is the usual inner structure algebra for a Jordan algebra described in Section 1.2.2, and the \mathbb{Z}_2 -grading produced by ε is $\mathfrak{Inst}(J, -)_{\bar{0}} = [R_J, R_J]$, $\mathfrak{Inst}(J, -)_{\bar{1}} = R_{J_0}$.

Example 1.10. If $(C_1, -)$ and $(C_2, -)$ are composition algebras over the field \mathbb{F} , then $(C_1 \otimes C_2, -)$ is a structurable algebra (see [6, Example 6.6]) for the product given by $(a \otimes b)(c \otimes d) = ac \otimes bd$ and the involution given by $\overline{a \otimes b} = \bar{a} \otimes \bar{b}$.

2 Models of the exceptional Lie algebras

The study of the Lie algebras began at the end of the XIX century, once Lie had translated certain problem of transformation groups to an algebraic context. The first fundamental contributions are due to Killing, who classified the complex simple Lie algebras in four key papers published during the years 1886 and 1890. Initially he thought that the only possible cases were the Lie algebras of the special linear group $SL_n(\mathbb{C})$ and of the orthogonal and symplectic groups $O(n, \mathbb{C})$ and $Sp(n, \mathbb{C})$, now called the classical Lie algebras. But, during his work, he obtained that, besides the classical Lie algebras, there were a few other Lie algebras, of dimensions 78, 133, 248, 52 and 14, now denoted as \mathfrak{e}_6 , \mathfrak{e}_7 , \mathfrak{e}_8 , \mathfrak{f}_4 and \mathfrak{g}_2 . Actually, he only proved the existence for \mathfrak{g}_2 , but he described all possibilities for rank, dimension and root systems. He found six algebras, since he did not notice that two of them were isomorphic (case \mathfrak{f}_4). This is a marvelous result, but the standard reference for it is Cartan's thesis in 1894, which completed Killing's classification, giving a rigorous treatment. This is a fundamental contribution, where Cartan proved the existence of all the exceptional simple Lie algebras.

2.1 First Models

The history of these algebras has been growing in parallel to the one of the related Lie groups. The first description of the smallest of the exceptional Lie groups was due to Engel ([25]), who, in 1900, described it as the isotropy group of a generic 3-form in 7 dimensions. Élie Cartan was the first to consider the group G_2 as the automorphism group of the octonion algebra in 1914 ([11, p. 298] although he commented about it earlier), as well the Lie algebra \mathfrak{g}_2 as the derivation algebra of the octonions (both on the split and compact forms). Jacobson generalized this result to arbitrary fields ([34]). This approach became popular through the article [26] by Hans Freudenthal, in 1951. But, for rather a long time, G_2 was the only Lie group for which further results were obtained.

The following model of a exceptional Lie algebra had to wait until 1950, when Chevalley and Schafer ([12]) showed that the set of derivations of the Albert algebra \mathbb{A} is \mathfrak{f}_4 . Tomber proved in [51] the converse: a Lie algebra over a field of characteristic 0 is of type F_4 if and only if it is isomorphic to the derivation algebra of an exceptional simple Jordan algebra. This fact led Tits, among other authors, to study the relationship between Jordan algebras and the remaining exceptional simple Lie algebras, which were constructed in a unified way [49]. We will revise this construction in the following subsection.

The algebra \mathfrak{e}_6 is also closely related to the Albert algebra. On one hand, it is the inner structure algebra of the Albert algebra (the Lie algebra generated by the right multiplication operators R_a for $a \in \mathbb{A}$ with zero trace, as

in Section 1.2.2). On the other hand, if $N: \mathbb{A} \rightarrow \mathbb{F}$ is the cubic norm as in Equation (2), and $N(a, b, c)$ denotes the trilinear form obtained by polarization, then the Lie algebra \mathfrak{e}_6 can be characterized as the Lie algebra $\{f \in \text{End}(\mathbb{A}) \mid N(f(a), b, c) + N(a, f(b), c) + N(a, b, f(c)) = 0\}$ ([35] and [26]).

The first model of \mathfrak{e}_7 related to the Albert algebra was provided in [50], as the algebra defined on the vector space $(\mathbb{A} \otimes \mathfrak{sl}_2(\mathbb{F})) \oplus \text{Der}(\mathbb{A})$ with the product

$$[x \otimes a + d_1, y \otimes b + d_2] = xy \otimes [a, b] + d_2(x) \otimes a - d_1(y) \otimes b - \frac{1}{2} \text{tr}(\text{ad } a \text{ ad } b)[R_x, R_y] + [d_1, d_2],$$

for $x, y \in \mathbb{A}$, $a, b \in \mathfrak{sl}_2(\mathbb{F})$ and $d_1, d_2 \in \text{Der}(\mathbb{A})$. The details of this construction appear in [37, §9]. This is a version of what nowadays is called the Tits-Kantor-Koecher construction applied to the Albert algebra. The name refers to several constructions which appeared almost simultaneously, and turned out to be essentially equivalent. In Koecher's construction [41], one forms $\mathbb{A} \oplus \bar{\mathbb{A}} \oplus \mathfrak{Str}(\mathbb{A})$, where $\bar{\mathbb{A}}$ is simply a copy of the vector space \mathbb{A} , with the anticommutative product given by $[x, y] = 0 = [\bar{x}, \bar{y}]$, $[x, \bar{y}] = 2R_{xy} + 2[R_y, R_x]$ if $x, y \in \mathbb{A}$, and $[L, x] = L(x)$, $[L, \bar{x}] = \bar{L}(x)$, if $L = R_x + \sum_i [R_{x_i}, R_{y_i}] \in \mathfrak{Str}(\mathbb{A})$, where $\bar{L} = -R_x + \sum_i [R_{x_i}, R_{y_i}]$. This construction will be generalized in Section 2.4. Similar to the situation for \mathfrak{e}_6 , the Lie algebra \mathfrak{e}_7 can be characterized too as the set of linear transformations of certain vector space M leaving invariant a quartic form [27]. Here, as a vector space, M is $\mathbb{A} \oplus \bar{\mathbb{A}} \oplus \mathbb{F} \oplus \bar{\mathbb{F}}$. This will play an important role in our description of the gradings with automorphisms of order 4 involved, in Sections 3.7 and 3.8.

Finally, the difficulty of finding a good model for \mathfrak{e}_8 (coordinate free, that is, not given by means of generators and relations obtained from the root system) is that the nontrivial representation of minimal dimension for \mathfrak{e}_8 is the adjoint representation, so there is no hope to embed \mathfrak{e}_8 as a subalgebra of $\mathfrak{gl}(V)$ for some vector space V of smaller dimension. However, some other 'linear models' can help in this purpose. Let V be a vector space of dimension 9, then we can construct \mathfrak{e}_8 as the vector space

$$\bigwedge^3 V^* \oplus \mathfrak{sl}(V) \oplus \bigwedge^3 V,$$

with Lie bracket as in [29, Exercise 22.21], based on the trilinear map given by the usual wedge product $\bigwedge^3 V \otimes \bigwedge^3 V \otimes \bigwedge^3 V \rightarrow \bigwedge^9 V \cong \mathbb{F}$.

We refer to [37, 48, 28, 26] for these and other algebraic constructions of the exceptional Lie algebras. We stress the reference [46], where many models appear: [46, Chapter 5, §1] is devoted to models of exceptional Lie algebras associated to a Cayley algebra (over arbitrary fields of characteristic zero, with several references to the reals), while [46, Chapter 5, §2] provides other models

based on gradings. This shows the interactions between nice models and certain gradings, and this philosophy is certainly present in Sections 3.9 and 3.10.

2.2 Tits construction

In 1966, Tits gave a unified construction of the exceptional simple Lie algebras (over fields of characteristic not two and three) in [49]. The construction used a couple of ingredients: an alternative algebra of degree 2 and a Jordan algebra of degree 3. In case these ingredients are chosen to be Hurwitz algebras and Jordan algebras of hermitian 3×3 matrices over Hurwitz algebras, Freudenthal's magic square [28] is obtained. We recall the construction in our concrete case.

Let C be a Hurwitz algebra over \mathbb{F} with norm q , and let $J = H_3(C', *)$ be the Jordan algebra of hermitian 3×3 -matrices over another Hurwitz algebra C' . Consider the vector space

$$\mathcal{T}(C, J) = \text{Der}(C) \oplus (C_0 \otimes J_0) \oplus \text{Der}(J)$$

with anticommutative multiplication specified by

- $\text{Der}(C)$ and $\text{Der}(J)$ are Lie subalgebras,
- $[\text{Der}(C), \text{Der}(J)] = 0$,
- $[d, a \otimes x] = d(a) \otimes x$, $[D, a \otimes x] = a \otimes D(x)$,
- $[a \otimes x, b \otimes y] = t_J(xy)d_{a,b} + [a, b] \otimes x * y + 2t_C(ab)[R_x, R_y]$

for all $d \in \text{Der}(C)$, $D \in \text{Der}(J)$, $a, b \in C_0$ and $x, y \in J_0$, with the notations of Sections 1.2.1 and 1.2.2. Now, using all the possibilities for C and C' , we obtain Freudenthal's Magic Square as follows [49] (note that we have added a column with $J = \mathbb{F}$ to obtain G_2 with the same construction):

		J				
$\mathcal{T}(C, J)$		\mathbb{F}	$H_3(\mathbb{F})$	$H_3(\mathbb{F} \times \mathbb{F})$	$H_3(\text{Mat}_{2 \times 2}(\mathbb{F}))$	$H_3(C)$
C	\mathbb{F}	0	\mathfrak{a}_1	\mathfrak{a}_2	\mathfrak{c}_3	\mathfrak{f}_4
	$\mathbb{F} \times \mathbb{F}$	0	\mathfrak{a}_2	$\mathfrak{a}_2 \oplus \mathfrak{a}_2$	\mathfrak{a}_5	\mathfrak{e}_6
	$\text{Mat}_{2 \times 2}(\mathbb{F})$	\mathfrak{a}_1	\mathfrak{c}_3	\mathfrak{a}_5	\mathfrak{d}_6	\mathfrak{e}_7
	C	\mathfrak{g}_2	\mathfrak{f}_4	\mathfrak{e}_6	\mathfrak{e}_7	\mathfrak{e}_8

2.3 Some symmetric constructions

In spite of the apparent asymmetry in the usage of the two Hurwitz algebras in Tits construction, the Magic Square is symmetric. This lead several authors to look for more symmetric constructions. Such an approach was taken by Vinberg in [46, p. 177], and interpreted by Barton and Sudbery in [10] as a construction depending on two composition algebras and closely related to the triality principle. A similar construction was provided by Landsberg and Manivel in [43], inspired by previous work of Allison and Faulkner [6].

The construction we are going to recall here (and use later on), is the construction in [18], based on two symmetric composition algebras, which has turned to be very useful in finding fine gradings on exceptional Lie algebras [19].

Let $(S, *, q)$ be a symmetric composition algebra and let

$$\mathfrak{o}(S, q) = \{d \in \text{End}_{\mathbb{F}}(S) \mid q(d(x), y) + q(x, d(y)) = 0 \forall x, y \in S\}$$

be the corresponding orthogonal Lie algebra. Consider the subalgebra of $\mathfrak{o}(S, q)^3$ defined by

$$\mathfrak{tri}(S, *, q) = \{(d_0, d_1, d_2) \in \mathfrak{o}(S, q)^3 \mid d_0(x * y) = d_1(x) * y + x * d_2(y) \forall x, y \in S\},$$

which is called the *triality Lie algebra*. The order three automorphism ϑ given by

$$\vartheta: \mathfrak{tri}(S, *, q) \longrightarrow \mathfrak{tri}(S, *, q), \quad (d_0, d_1, d_2) \longmapsto (d_2, d_0, d_1),$$

is called the *triality automorphism*. Take the element of $\mathfrak{tri}(S, *, q)$ (denoted by $\mathfrak{tri}(S)$ when it is no ambiguity) given by

$$t_{x,y} := \left(\sigma_{x,y}, \frac{1}{2}q(x, y)id - r_x l_y, \frac{1}{2}q(x, y)id - l_x r_y \right),$$

where $\sigma_{x,y}(z) = q(x, z)y - q(y, z)x$, $r_x(z) = z * x$, and $l_x(z) = x * z$ for any $x, y, z \in S$.

Let $(S, *, q)$ and (S', \star, q') be two symmetric composition algebras over \mathbb{F} . Consider the following vector space, which depends symmetrically on S and S' :

$$\mathfrak{g}(S, S') := \mathfrak{tri}(S, *, q) \oplus \mathfrak{tri}(S', \star, q') \oplus \left(\bigoplus_{i=0}^2 \iota_i(S \otimes S') \right)$$

where $\iota_i(S \otimes S')$ is just a copy of $S \otimes S'$ ($i = 0, 1, 2$), and the anticommutative product on $\mathfrak{g}(S, S')$ is determined by the following conditions:

- $\mathfrak{tri}(S, *, q) \oplus \mathfrak{tri}(S', \star, q')$ is a Lie subalgebra of $\mathfrak{g}(S, S')$;

- $[(d_0, d_1, d_2), \iota_i(x \otimes x')] = \iota_i(d_i(x) \otimes x')$, $[(d'_0, d'_1, d'_2), \iota_i(x \otimes x')] = \iota_i(x \otimes d'_i(x'))$, for any $(d_0, d_1, d_2) \in \mathbf{tri}(S)$, $(d'_0, d'_1, d'_2) \in \mathbf{tri}(S')$, $x \in S$ and $x' \in S'$;
- $[\iota_i(x \otimes x'), \iota_{i+1}(y \otimes y')] = \iota_{i+2}((x * y) \otimes (x' * y'))$ (indices modulo 3), for any $i = 0, 1, 2$, $x, y \in S$ and $x', y' \in S'$;
- $[\iota_i(x \otimes x'), \iota_i(y \otimes y')] = q'(x', y')\vartheta^i(t_{x,y}) + q(x, y)\vartheta'^i(t'_{x',y'}) \in \mathbf{tri}(S) \oplus \mathbf{tri}(S')$, for any $i = 0, 1, 2$, $x, y \in S$ and $x', y' \in S'$, ϑ and ϑ' being the corresponding triality automorphisms.

The anticommutative algebra $\mathfrak{g}(S, S')$ defined in this way turns out to be a Lie algebra ([18, Theorem 3.1]), and we recover Freudenthal's Magic Square if symmetric composition algebras of all possible dimensions are considered:

		dim S			
		1	2	4	8
dim S'	1	\mathfrak{a}_1	\mathfrak{a}_2	\mathfrak{c}_3	\mathfrak{f}_4
	2	\mathfrak{a}_2	$\mathfrak{a}_2 \oplus \mathfrak{a}_2$	\mathfrak{a}_5	\mathfrak{e}_6
	4	\mathfrak{c}_3	\mathfrak{a}_5	\mathfrak{d}_6	\mathfrak{e}_7
	8	\mathfrak{f}_4	\mathfrak{e}_6	\mathfrak{e}_7	\mathfrak{e}_8

If C_1 and C_2 are two Hurwitz algebras over \mathbb{F} , the Lie algebra $\mathfrak{g}(pC_1, pC_2)$ is isomorphic to $\mathcal{T}(C_1, H_3(C_2, *))$ [23, Theorem 6.25].

2.4 Kantor's construction

Let $(A, -)$ be a structurable algebra. Denote by $S \equiv S(A, -)$ its set of skew-hermitian elements. Endow the \mathbb{Z} -graded vector space $\mathcal{K} = \mathcal{K}_{-2} \oplus \mathcal{K}_{-1} \oplus \mathcal{K}_0 \oplus \mathcal{K}_1 \oplus \mathcal{K}_2$, for

$$\begin{aligned} \mathcal{K}_2 &= S, & \mathcal{K}_1 &= A, & \mathcal{K}_0 &= \mathfrak{Inst}(A, -), \\ \mathcal{K}_{-2} &= S^\sim & \mathcal{K}_{-1} &= A^\sim, \end{aligned}$$

where S^\sim and A^\sim are simply copies of S and A respectively, with a graded Lie algebra structure given by the anticommutative multiplication such that $\mathfrak{Inst}(A, -) = V_{A,A}$ is a subalgebra and the following conditions hold:

$$\begin{aligned} [T, a] &= T(a), & [T, a^\sim] &= (T^\varepsilon a)^\sim, \\ [T, s] &= T(s) + s\overline{T(1)}, & [T, s^\sim] &= (T^\varepsilon(s) + s\overline{T^\varepsilon(1)})^\sim, \\ [a + s, a' + s'] &= 2(s\bar{s}' - s'\bar{s}) \in \mathcal{K}_2, \\ [a^\sim + s^\sim, a'^\sim + s'^\sim] &= 2(\overline{s\bar{s}'} - \overline{s'\bar{s}})^\sim \in \mathcal{K}_{-2}, \\ [a + s, a'^\sim + s'^\sim] &= (-s'a)^\sim + L_s L_{s'} + 2V_{a,a'} + (sa') \in \mathcal{K}_{-1} \oplus \mathcal{K}_0 \oplus \mathcal{K}_1, \end{aligned} \tag{4}$$

for $T \in \mathfrak{Inst}(A, -)$, $a, a' \in A$, $s, s' \in S$, where $L_s: A \rightarrow A$ denotes the left multiplication by $s \in S$ (so that $2L_s L_{s'} = V_{s s', 1} - V_{s, s'} \in \mathfrak{Inst}(A, -)$). Following

[4, §6.4], this is a \mathbb{Z} -graded Lie algebra denoted by $\mathfrak{Kan}(A, -)$, which is called the *Kantor construction* attached to the structurable algebra A . The construction takes its name from Kantor, who introduced it first in [40], although in a somewhat different way.

In connection with Example 1.9, if J is a Jordan algebra, the Kantor construction $\mathfrak{Kan}(J, -)$ coincides (up to isomorphism) with the classical Tits-Kantor-Koecher Lie algebra constructed from the Jordan algebra J .

A necessary and sufficient condition for a Lie algebra to be isomorphic to the Kantor's construction attached to a structurable algebra is given by the existence of an \mathfrak{sl}_2 -triple $\{e, h, f\}$ (that is, $[h, e] = 2e, [h, f] = -2f, [e, f] = h$) in L such that L is the direct sum of irreducible modules for $\langle \{e, h, f\} \rangle$ of dimensions 1, 3, and 5; the only ideal of L which centralizes $\{e, h, f\}$ is $\{0\}$, and L is generated by the eigenspaces 2 and -2 for $\text{ad } h$ [4, Theorem 6.10].

The relevance for our purposes comes from the fact that exceptional Lie algebras are obtained in terms of Kantor's constructions attached to certain structurable algebras. Some examples of this situation are shown next:

Example 2.1. Consider the tensor products of composition algebras: $\mathcal{C} = \mathbb{F} \otimes \mathcal{C}$, $\mathcal{K} \otimes \mathcal{C}$, $\mathcal{Q} \otimes \mathcal{C}$, and $\mathcal{C} \otimes \mathcal{C}$. These are structurable algebras according to Example 1.10. Kantor's construction gives the following Lie algebras:

$$\mathfrak{Kan}(\mathcal{C}) \cong \mathfrak{f}_4, \quad \mathfrak{Kan}(\mathcal{K} \otimes \mathcal{C}) \cong \mathfrak{e}_6, \quad \mathfrak{Kan}(\mathcal{Q} \otimes \mathcal{C}) \cong \mathfrak{e}_7, \quad \mathfrak{Kan}(\mathcal{C} \otimes \mathcal{C}) \cong \mathfrak{e}_8,$$

as stated in [40] (see, alternatively, [3, §8(c)]).

Example 2.2. Let $J = H_4(C, *)$ be the Jordan algebra defined in Section 1.2.2 for C a Hurwitz associative algebra. This algebra is a finite-dimensional simple Jordan algebra, with generic trace $t_J(x) = \frac{1}{4} \text{tr}(x)$. We can use a sort of Cayley-Dickson process with J in order to construct a structurable algebra as follows. Take a nonzero $\mu \in \mathbb{F}$ and the algebra defined on the vector space

$$A = J \oplus vJ,$$

where v is simply a convenient mark to indicate the cartesian product of two copies of J , with multiplication and involution given by

$$\begin{aligned} (x_1 + vx_2)(x_3 + vx_4) &:= x_1x_3 + \mu(x_2x_4^\vartheta)^\vartheta + v(x_1^\vartheta x_4 + (x_2^\vartheta x_3^\vartheta)^\vartheta), \\ \overline{x_1 + vx_2} &:= x_1 - vx_2^\vartheta \end{aligned} \tag{5}$$

for any $x_i \in J$ for $i = 1, 2, 3, 4$, where the multiplication in J is denoted by juxtaposition, with $x^\vartheta := -x + 2t_J(x)1$ (so that $1^\vartheta = 1$) for any $x \in J$. The resulting algebra $(A, -)$ is denoted by $CD(H_4(C))$. According to [5] (see, alternatively, [4, Example 6.7]), this algebra is structurable with the extra property that the

space of skew-hermitian elements has dimension 1. The Lie algebra obtained by means of Kantor's construction $\mathfrak{Kan}(CD(H_4(C)))$ is \mathfrak{e}_6 , \mathfrak{e}_7 and \mathfrak{e}_8 , respectively, in case $\dim C = 1, 2$ or 4 .

2.5 Steinberg's construction

Consider the following example [6].

Example 2.3. Let $(A, -)$ be a unital associative algebra with involution and consider the unitary Lie algebra $\mathfrak{u}_n(A, -) = \{x \in \text{Mat}_{n \times n}(A) \mid \bar{x}^t = -x\}$. Some remarkable elements are $u_{ij}(a) = ae_{ij} - \bar{a}e_{ji}$, where e_{ij} are the usual matrix units. These elements are subject to the following relations:

$$\begin{aligned} u_{ij}(a) &= u_{ji}(-\bar{a}), \\ a &\mapsto u_{ij}(a) \text{ is a linear map,} \\ [u_{ij}(a), u_{jk}(b)] &= u_{ik}(ab) \text{ for distinct } i, j, k, \\ [u_{ij}(a), u_{kl}(b)] &= 0 \text{ for distinct } i, j, k, l. \end{aligned} \tag{6}$$

Now, let $(A, -)$ be a unital algebra with involution, and let $\mathfrak{stu}_n(A, -)$ denote the Lie subalgebra generated by elements $u_{ij}(a)$, for $1 \leq i \neq j \leq n$, $n \geq 3$, and $a \in A$, subject to the relations (6). Then the condition $u_{ij}(a) = 0$ implies $a = 0$ if and only if either $n \geq 4$ and A is associative or $n = 3$ and $(A, -)$ is structurable. This Lie algebra is called the *Steinberg unitary Lie algebra* by analogy with the Steinberg group in K-theory.

During the proof of the previous result, Allison and Faulkner used the following construction (also developed in [6]).

Let $(A, -)$ be a structurable algebra. For $T \in \mathfrak{gl}(A)$, define \bar{T} by $\bar{T}(x) = \overline{T(\bar{x})}$. A set $T = (T_1, T_2, T_3) \in \mathfrak{gl}(A)^3$ is said to be a *related triple* if

$$\bar{T}_i(xy) = T_{i+1}(x)y + xT_{i+2}(y)$$

for all $x, y \in A$ and for all $i = 1, 2, 3$, where the subindices are taken modulo 3. These triples form a Lie algebra denoted by $\mathfrak{trip}(A)$. A remarkable triple is the following:

$$\begin{aligned} T_i &= L_{\bar{b}}L_a - L_{\bar{a}}L_b, \\ T_{i+1} &= R_{\bar{b}}R_a - R_{\bar{a}}R_b, \\ T_{i+2} &= R_{\bar{a}b-\bar{b}a} + L_bL_{\bar{a}} - L_aL_{\bar{b}}, \end{aligned} \tag{7}$$

for $a, b \in A$, where L_a and R_a denote the left and right multiplications by a in A .

Consider now the vector space

$$\mathcal{U}(A, -) := \mathfrak{trip}(A) \oplus u_{12}(A) \oplus u_{23}(A) \oplus u_{31}(A)$$

with anticommutative multiplication such that $\mathbf{trip}(A)$ is a subalgebra and the following conditions hold:

$$\begin{aligned} [T, u_{i,i+1}(a)] &= u_{i,i+1}(T_{i+2}(a)), \\ [u_{i,i+1}(a), u_{i+1,i+2}(b)] &= -u_{i+2,i}(\overline{ab}), \\ [u_{i,i+1}(a), u_{i,i+1}(b)] &= T \text{ as in (7),} \end{aligned}$$

for any $a, b \in A$ and $T = (T_1, T_2, T_3) \in \mathbf{trip}(A)$. Then $\mathcal{U}(A, -)$ is a Lie algebra isomorphic to $\mathfrak{stu}_3(A, -)/\mathfrak{z}$, where \mathfrak{z} is the center of $\mathfrak{stu}_3(A, -)$ ([6, Theorem 4.3]). This algebra $\mathcal{U}(A, -)$ is simple if and only if $(A, -)$ is so. We call this algebra the *Steinberg construction* attached to the structurable algebra A . Moreover, it turns out that $\mathcal{U}(A, -)$ is isomorphic to the Lie algebra given by the Kantor's construction $\mathfrak{Kan}(A, -)$ considered in the previous subsection.

3 Description of the gradings

3.1 Gradings on Hurwitz algebras

Given a Hurwitz algebra C with norm q , we can construct, for each $0 \neq \alpha \in \mathbb{F}$, a new unital algebra with involution, denoted by $\mathfrak{CD}(C, \alpha)$, by means of the so called Cayley-Dickson doubling process. This is the algebra defined on $C \times C$ with the multiplication given by

$$(a, b)(c, d) = (ac + \alpha \bar{d}b, da + b\bar{c})$$

and the quadratic form

$$q(a, b) = q(a) - \alpha q(b).$$

The resulting algebra $\mathfrak{CD}(C, \alpha)$ is a Hurwitz algebra if and only if C is associative. Thus, according to our list of Hurwitz algebras, $\mathfrak{CD}(\mathbb{F}, \alpha)$ is isomorphic to \mathcal{K} , $\mathfrak{CD}(\mathcal{K}, \alpha)$ is isomorphic to \mathcal{Q} and $\mathfrak{CD}(\mathcal{Q}, \alpha)$ is isomorphic to \mathcal{C} . The algebra $\mathfrak{CD}(C, \alpha)$ is always \mathbb{Z}_2 -graded, with even part $\{(a, 0) \mid a \in C\}$ and odd part $\{(0, b) \mid b \in C\}$. In particular \mathcal{K} is \mathbb{Z}_2 -graded, \mathcal{Q} is \mathbb{Z}_2^2 -graded and \mathcal{C} is \mathbb{Z}_2^3 -graded.

On the other hand the standard basis of \mathcal{C} is associated to a fine \mathbb{Z}^2 -grading, called the *Cartan grading* on \mathcal{C} , which is given by

$$\begin{aligned} \mathcal{C}_{(0,0)} &= \mathbb{F}e_1 \oplus \mathbb{F}e_2, \\ \mathcal{C}_{(1,0)} &= \mathbb{F}u_1, & \mathcal{C}_{(-1,0)} &= \mathbb{F}v_1, \\ \mathcal{C}_{(0,1)} &= \mathbb{F}u_2, & \mathcal{C}_{(0,-1)} &= \mathbb{F}v_2, \\ \mathcal{C}_{(-1,-1)} &= \mathbb{F}u_3, & \mathcal{C}_{(1,1)} &= \mathbb{F}v_3. \end{aligned} \tag{8}$$

The subalgebra $\langle e_1, e_2, u_1, v_1 \rangle$ can be identified with \mathcal{Q} , which turns out to be \mathbb{Z} -graded, for

$$\mathcal{Q}_0 = \mathbb{F}e_1 \oplus \mathbb{F}e_2, \quad \mathcal{Q}_1 = \mathbb{F}u_1, \quad \mathcal{Q}_{-1} = \mathbb{F}v_1. \quad (9)$$

3.2 Gradings on symmetric composition algebras

Gradings on symmetric composition algebras were classified in [19, Theorem 4.5]. Every group grading on a Hurwitz algebra C is a grading on pC (since the involution preserves the homogeneous components), and the gradings on both the Hurwitz algebra C and its para-Hurwitz counterpart coincide when C has dimension at least 4. In particular we have a \mathbb{Z}_2 -grading on $p\mathcal{K}$, a \mathbb{Z}_2^2 -grading and a \mathbb{Z} -grading on $p\mathcal{Q}$ and a \mathbb{Z}_3^3 -grading and a \mathbb{Z}^2 -grading on $p\mathcal{C}$.

There is a remarkable \mathbb{Z}_3 -grading in the case of dimension 2 which does not come from a grading on the corresponding Hurwitz algebra, namely,

$$p\mathcal{K}_{\bar{0}} = 0, \quad p\mathcal{K}_{\bar{1}} = \mathbb{F}e_1 \quad p\mathcal{K}_{\bar{2}} = \mathbb{F}e_2,$$

where e_1 and e_2 are the orthogonal idempotents $(1, 0)$ and $(0, 1)$ in $\mathcal{K} = \mathbb{F} \times \mathbb{F}$, which in $p\mathcal{K}$ satisfy $e_1 * e_1 = e_2$ and $e_2 * e_2 = e_1$.

Also, a natural \mathbb{Z}_3^2 -grading appears on the pseudo-octonion algebra $\text{Ok} = (\mathfrak{sl}_3(\mathbb{F}), *, q)$, determined by

$$\text{Ok}_{(\bar{1}, \bar{0})} = \mathbb{F} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{pmatrix} \quad \text{and} \quad \text{Ok}_{(\bar{0}, \bar{1})} = \mathbb{F} \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$$

(recall that ω is a primitive cubic root of 1).

3.3 Gradings on Lie algebras induced from gradings on symmetric composition algebras

If S and S' are two symmetric composition algebras, the Lie algebra $\mathfrak{g}(S, S')$ is always \mathbb{Z}_2^2 -graded, for

$$\begin{aligned} \mathfrak{g}(S, S')_{(0,0)} &= \text{tri}(S) \oplus \text{tri}(S'), & \mathfrak{g}(S, S')_{(1,1)} &= \iota_0(S \otimes S'), \\ \mathfrak{g}(S, S')_{(1,0)} &= \iota_1(S \otimes S'), & \mathfrak{g}(S, S')_{(0,1)} &= \iota_2(S \otimes S'). \end{aligned} \quad (10)$$

Moreover, if $S = \bigoplus_{g \in G} S_g$ is G -graded, and $S' = \bigoplus_{g \in G'} S'_g$ is G' -graded, these gradings can be combined with the one in Equation (10), thus obtaining a grading on $\mathfrak{g}(S, S')$ by the group $\mathbb{Z}_2^2 \times G \times G'$.

As a consequence, if we consider the gradings by the groups \mathbb{Z}_2^r on the symmetric composition algebras as in Section 3.2, we get gradings on:

(6g1) $\epsilon_6 = \mathfrak{g}(p\mathcal{K}, p\mathcal{C})$ by the group $\mathbb{Z}_2^2 \times \mathbb{Z}_2 \times \mathbb{Z}_2^3 = \mathbb{Z}_2^6$, of type (48, 1, 0, 7);

(7g1) $\epsilon_7 = \mathfrak{g}(p\mathcal{Q}, p\mathcal{C})$ by the group $\mathbb{Z}_2^2 \times \mathbb{Z}_2^2 \times \mathbb{Z}_2^3 = \mathbb{Z}_2^7$, of type (96, 0, 3, 7);

(8g1) $\epsilon_8 = \mathfrak{g}(p\mathcal{C}, p\mathcal{C})$ by the group $\mathbb{Z}_2^2 \times \mathbb{Z}_2^3 \times \mathbb{Z}_2^3 = \mathbb{Z}_2^8$, of type (192, 0, 0, 14).

Moreover, there is a distinguished \mathbb{Z}_3 -grading on $\mathfrak{g}(S, S')$, obtained as the eigenspace decomposition of an order 3 automorphism Θ induced by the triality automorphisms ϑ and ϑ' of S and S' respectively. It is given by

$$\Theta|_{\text{tri}(S)} = \vartheta, \quad \Theta|_{\text{tri}(S')} = \vartheta', \quad \Theta(\iota_i(x \otimes x')) = \iota_{i+1}(x \otimes x').$$

This \mathbb{Z}_3 -grading on $\mathfrak{g}(S, S')$ induced by Θ can be combined with any G and G' -gradings on S and S' respectively, to get a grading by the group $\mathbb{Z}_3 \times G \times G'$ on $\mathfrak{g}(S, S')$. If this procedure is applied to the \mathbb{Z}_3 -grading on $p\mathcal{K}$ and to the \mathbb{Z}_3^2 -grading on Ok , we get gradings on:

(6g2) $\epsilon_6 = \mathfrak{g}(p\mathcal{K}, \text{Ok})$ by the group $\mathbb{Z}_3 \times \mathbb{Z}_3 \times \mathbb{Z}_3^2 = \mathbb{Z}_3^4$, of type (72, 0, 2);

(8g2) $\epsilon_8 = \mathfrak{g}(\text{Ok}, \text{Ok})$ by the group $\mathbb{Z}_3 \times \mathbb{Z}_3^2 \times \mathbb{Z}_3^2 = \mathbb{Z}_3^5$, of type (240, 0, 0, 2).

And if we combine some of the gradings above (related to the primes 2 and 3), we obtain gradings on:

(6g3) $\epsilon_6 = \mathfrak{g}(p\mathcal{K}, p\mathcal{C})$ by the group $\mathbb{Z}_3 \times \mathbb{Z}_3 \times \mathbb{Z}_2^3 = \mathbb{Z}_2^3 \times \mathbb{Z}_3^2$, of type (64, 7);

(6g4) $\epsilon_6 = \mathfrak{g}(p\mathcal{K}, \text{Ok})$ by the group $\mathbb{Z}_3 \times \mathbb{Z}_2 \times \mathbb{Z}_3^2 = \mathbb{Z}_2 \times \mathbb{Z}_3^3$, of type (26, 26);

(7g2) $\epsilon_7 = \mathfrak{g}(p\mathcal{Q}, \text{Ok})$ by the group $\mathbb{Z}_3 \times \mathbb{Z}_2^2 \times \mathbb{Z}_3^2 = \mathbb{Z}_2^2 \times \mathbb{Z}_3^3$, of type (81, 26);

(8g3) $\epsilon_8 = \mathfrak{g}(p\mathcal{C}, \text{Ok})$ by the group $\mathbb{Z}_3 \times \mathbb{Z}_2^3 \times \mathbb{Z}_3^2 = \mathbb{Z}_2^3 \times \mathbb{Z}_3^3 = \mathbb{Z}_6^3$, of type (182, 33).

Finally, we can also combine the above \mathbb{Z}_3 -grading on $\mathfrak{g}(S, S')$ induced by Θ with gradings on S and S' by infinite groups, namely, the \mathbb{Z} -grading on $p\mathcal{Q}$ and the \mathbb{Z}^2 -grading on $p\mathcal{C}$ as in Equations (9) and (8) respectively. Thus we get gradings on:

(6g5) $\epsilon_6 = \mathfrak{g}(p\mathcal{K}, p\mathcal{C})$ by the group $\mathbb{Z}_3 \times \mathbb{Z}_3 \times \mathbb{Z}^2 = \mathbb{Z}^2 \times \mathbb{Z}_3^2$, of type (60, 9);

(7g3) $\epsilon_7 = \mathfrak{g}(p\mathcal{Q}, \text{Ok})$ by the group $\mathbb{Z}_3 \times \mathbb{Z} \times \mathbb{Z}_3^2 = \mathbb{Z} \times \mathbb{Z}_3^3$, of type (55, 0, 26);

(8g4) $\epsilon_8 = \mathfrak{g}(p\mathcal{C}, \text{Ok})$ by the group $\mathbb{Z}_3 \times \mathbb{Z}^2 \times \mathbb{Z}_3^2 = \mathbb{Z}^2 \times \mathbb{Z}_3^3$, of type (168, 1, 26).

Now, consider the \mathbb{Z} -grading on $\mathfrak{g}(S, S')$ described in [7, §4.2] as follows: Let 1 and 1' be the unit elements of two Hurwitz algebras C and C' . They become the paraunits of the corresponding para-Hurwitz algebras $S = pC$ and $S' = pC'$. Consider the inner derivation $d := \text{ad}(\iota_0(1 \otimes 1'))$ of $\mathfrak{g}(S, S')$, which is a semisimple derivation with eigenvalues $\pm 2, \pm 1, 0$. Thus, the eigenspace decomposition gives the following \mathbb{Z} -grading (5-grading) on $\mathfrak{g}(S, S')$:

$$\begin{aligned} \mathfrak{g}(S, S')_{\pm 2} &= \Sigma_{\pm}(S_0, S'_0), \\ \mathfrak{g}(S, S')_{\pm 1} &= \nu_{\pm}(S \otimes S'), \\ \mathfrak{g}(S, S')_0 &= t_{S_0, S_0} \oplus t_{S'_0, S'_0} \oplus \iota_0(S_0 \otimes S'_0) \oplus \mathbb{F}\iota_0(1 \otimes 1'), \end{aligned} \quad (11)$$

where S_0 and S'_0 denote the subspaces of zero trace elements in C and C' (here $S = pC$ and $S' = pC'$), and where

$$\begin{aligned} \Sigma_{\pm}(y, y') &:= t_{1,y} + t'_{1,y'} \pm \mathbf{i}\iota_0(y \otimes 1 + 1 \otimes y'), \\ \nu_{\pm}(y \otimes y') &:= \iota_1(y \otimes y') \mp \mathbf{i}\iota_2(\bar{y} \otimes \bar{y}'), \end{aligned}$$

for all $y \in S$, $y' \in S'$, and for a fixed scalar $\mathbf{i} \in \mathbb{F}$ such that $\mathbf{i}^2 = -1$. It is clear that this \mathbb{Z} -grading can be refined with gradings coming from S or S' . In particular, when the \mathbb{Z}_2^r -gradings on pC and pC' are used, we get gradings on:

- (6g6) $\mathfrak{e}_6 = \mathfrak{g}(p\mathcal{K}, pC)$ by the group $\mathbb{Z} \times \mathbb{Z}_2 \times \mathbb{Z}_2^3 = \mathbb{Z} \times \mathbb{Z}_2^4$, of type (57, 0, 7);
- (7g4) $\mathfrak{e}_7 = \mathfrak{g}(p\mathcal{Q}, pC)$ by the group $\mathbb{Z} \times \mathbb{Z}_2^2 \times \mathbb{Z}_2^3 = \mathbb{Z} \times \mathbb{Z}_2^5$, of type (106, 3, 7);
- (8g5) $\mathfrak{e}_8 = \mathfrak{g}(pC, pC)$ by the group $\mathbb{Z} \times \mathbb{Z}_2^3 \times \mathbb{Z}_2^3 = \mathbb{Z} \times \mathbb{Z}_2^6$, of type (206, 0, 14).

Next we recall the \mathbb{Z}^4 -grading on $\mathfrak{g}(pC, S')$ described in [7, §4.3]. Take the canonical generators $a_1 = (1, 0, 0, 0)$, $a_2 = (0, 1, 0, 0)$, $g_1 = (0, 0, 1, 0)$, $g_2 = (0, 0, 0, 1)$ of the group \mathbb{Z}^4 , and write $a_0 = -a_1 - a_2$, $g_0 = -g_1 - g_2$. Set the degrees of the \mathbb{Z}^4 -grading as follows:

$$\begin{aligned} \deg \iota_i(e_1 \otimes s) &= a_i &= -\deg \iota_i(e_2 \otimes s), \\ \deg \iota_i(u_i \otimes s) &= g_i &= -\deg \iota_i(v_i \otimes s), \\ \deg \iota_i(u_{i+1} \otimes s) &= a_{i+2} + g_{i+1} &= -\deg \iota_i(v_{i+1} \otimes s), \\ \deg \iota_i(u_{i+2} \otimes s) &= -a_{i+1} + g_{i+2} &= -\deg \iota_i(v_{i+2} \otimes s), \end{aligned} \quad (12)$$

where $s \in S'$ and $B = \{e_1, e_2, u_0, u_1, u_2, v_0, v_1, v_2\}$ is the standard basis of the algebra \mathcal{C} described in Section 1.2.1. Also set $\deg(t) = (0, 0, 0, 0)$ for all $t \in \text{tri}(S')$, and $\deg t_{x,y} = \deg \iota_0(x \otimes s) + \deg \iota_0(y \otimes s)$, if $x, y \in B$. A straightforward computation shows that the \mathbb{Z}^4 -grading on $\mathfrak{g}(pC, S')$ provided by this assignment is compatible with any grading on pC and on S' . In particular, the following gradings are obtained:

(6g7) A grading on $\mathfrak{e}_6 = \mathfrak{g}(p\mathcal{C}, p\mathcal{K})$ by the group $\mathbb{Z}^4 \times \mathbb{Z}_2$, of type (72, 1, 0, 1);

(7g5) A grading on $\mathfrak{e}_7 = \mathfrak{g}(p\mathcal{C}, p\mathcal{Q})$ by the group $\mathbb{Z}^4 \times \mathbb{Z}_2^2$, of type (120, 0, 3, 1);

(8g6) A grading on $\mathfrak{e}_8 = \mathfrak{g}(p\mathcal{C}, p\mathcal{C})$ by the group $\mathbb{Z}^4 \times \mathbb{Z}_2^3$, of type (216, 0, 0, 8).

Observe that a \mathbb{Z}^3 -grading can be defined on $\mathfrak{g}(p\mathcal{Q}, S')$ and a \mathbb{Z}^2 -grading on $\mathfrak{g}(p\mathcal{K}, S')$, both of them inherited directly from the \mathbb{Z}^4 -grading on $\mathfrak{g}(p\mathcal{C}, S')$ given by Equation (12). (Here $\{e_1, e_2, u_1, v_1\}$ is a basis of $p\mathcal{Q}$ and $\{e_1, e_2\}$ is a basis of $p\mathcal{K}$.) The grading on $\mathfrak{g}(p\mathcal{Q}, S')$ is given by the following assignment of degrees:

$$\begin{aligned} \deg \iota_i(e_1 \otimes s) &= a_i &= -\deg \iota_i(e_2 \otimes s), \\ \deg \iota_1(u_1 \otimes s) &= (0, 0, 1) &= -\deg \iota_1(v_1 \otimes s), \\ \deg \iota_2(u_1 \otimes s) &= (0, 1, 1) &= -\deg \iota_2(v_1 \otimes s), \\ \deg \iota_0(u_1 \otimes s) &= (1, 1, 1) &= -\deg \iota_0(v_1 \otimes s), \\ \deg(t_{e_1, e_2}) &= (0, 0, 0) &= \deg(t_{u_1, v_1}), \\ \deg(t_{e_1, u_1}) &= (-1, 0, 1) &= -\deg(t_{e_2, v_1}), \\ \deg(t_{e_1, v_1}) &= (-1, -2, -1) &= -\deg(t_{e_2, u_1}), \end{aligned}$$

for $a_1 = (1, 0, 0)$, $a_2 = (0, 1, 0)$, $a_0 = (-1, -1, 0)$, with $\deg(\mathbf{tri}(S')) = (0, 0, 0)$. The grading on $\mathfrak{g}(p\mathcal{K}, S')$ is given by:

$$\begin{aligned} \deg \iota_1(e_1 \otimes s) &= (1, 0) &= -\deg \iota_1(e_2 \otimes s), \\ \deg \iota_2(e_1 \otimes s) &= (0, 1) &= -\deg \iota_2(e_2 \otimes s), \\ \deg \iota_0(e_1 \otimes s) &= (-1, -1) &= -\deg \iota_0(e_2 \otimes s), \end{aligned}$$

with $\deg(\mathbf{tri}(p\mathcal{K})) = (0, 0) = \deg(\mathbf{tri}(S'))$. Again these gradings can be combined with the \mathbb{Z}_2^3 -grading on the symmetric composition algebra $p\mathcal{C}$ to get:

(6g8) A grading on $\mathfrak{e}_6 = \mathfrak{g}(p\mathcal{K}, p\mathcal{C})$ by the group $\mathbb{Z}^2 \times \mathbb{Z}_2^3$, of type (48, 1, 0, 7);

(7g6) A grading on $\mathfrak{e}_7 = \mathfrak{g}(p\mathcal{Q}, p\mathcal{C})$ by the group $\mathbb{Z}^3 \times \mathbb{Z}_2^3$, of type (102, 0, 1, 7).

3.4 Gradings on some Jordan algebras

We are going to describe some gradings on the Jordan algebra $H_4(C, *) = \{x = (x_{ij}) \in \text{Mat}_{4 \times 4}(C) \mid x_{ij} = \bar{x}_{ji}\}$ for C some associative Hurwitz algebra (that is, up to isomorphism, $C \in \{\mathbb{F}, \mathcal{K}, \mathcal{Q}\}$). The reader may consult [23, Chapter 5] for the description of gradings on simple Jordan algebras.

Observe first that the Kronecker product gives an isomorphism of associative algebras:

$$\text{Mat}_{2 \times 2}(\mathbb{F}) \otimes \text{Mat}_{2 \times 2}(\mathbb{F}) \rightarrow \text{Mat}_{4 \times 4}(\mathbb{F}), \quad a \otimes b \mapsto a \otimes b = \begin{pmatrix} a_{11}b & a_{12}b \\ a_{21}b & a_{22}b \end{pmatrix}.$$

Also, there is the natural isomorphism of associative algebras,

$$\mathrm{Mat}_{4 \times 4}(\mathbb{F}) \otimes C \cong \mathrm{Mat}_{4 \times 4}(C), \quad (a_{ij}) \otimes x \mapsto (a_{ij}x). \quad (13)$$

As $\mathrm{Mat}_{2 \times 2}(\mathbb{F})$ is isomorphic to \mathcal{Q} , it inherits a \mathbb{Z} -grading and a \mathbb{Z}_2^2 -grading, so that the previous identifications allow us to define gradings on $\mathrm{Mat}_{4 \times 4}(\mathbb{F})$ by the groups $\mathbb{Z}_2^2 \times \mathbb{Z}_2^2$ and $\mathbb{Z} \times \mathbb{Z}_2^2$, on $\mathrm{Mat}_{4 \times 4}(\mathcal{K})$ by the groups $\mathbb{Z}_2^2 \times \mathbb{Z}_2^2 \times \mathbb{Z}_2$ and $\mathbb{Z} \times \mathbb{Z}_2^2 \times \mathbb{Z}_2$, and on $\mathrm{Mat}_{4 \times 4}(\mathcal{Q})$ by the groups $\mathbb{Z}_2^2 \times \mathbb{Z}_2^2 \times \mathbb{Z}_2^2$ and $\mathbb{Z} \times \mathbb{Z}_2^2 \times \mathbb{Z}_2^2$. Trivially any grading on the associative algebra $\mathrm{Mat}_{4 \times 4}(C)$ is a grading of the Jordan algebra $\mathrm{Mat}_{4 \times 4}(C)^+$. The point is that, for the previously described gradings, the Jordan subalgebra $H_4(C, *) \leq \mathrm{Mat}_{4 \times 4}(C)^+$ is a graded subspace, so that:

- $H_4(\mathbb{F}, *)$ is \mathbb{Z}_2^4 and $\mathbb{Z} \times \mathbb{Z}_2^2$ -graded;
- $H_4(\mathcal{K}, *)$ is \mathbb{Z}_2^5 and $\mathbb{Z} \times \mathbb{Z}_2^3$ -graded;
- $H_4(\mathcal{Q}, *)$ is \mathbb{Z}_2^6 and $\mathbb{Z} \times \mathbb{Z}_2^4$ -graded.

Let us explain this with some extra detail. Let us denote by q_0 the identity matrix of degree 2, and consider the matrices

$$q_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad q_2 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad q_3 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} = q_1 q_2. \quad (14)$$

Then the assignment $\deg(q_1) = (\bar{1}, \bar{0})$ and $\deg(q_2) = (\bar{0}, \bar{1})$ gives the \mathbb{Z}_2^2 -grading on $\mathrm{Mat}_{2 \times 2}(\mathbb{F})$. The \mathbb{Z}_2^4 -grading on $\mathrm{Mat}_{4 \times 4}(\mathbb{F})$ has 16 one-dimensional homogeneous components, where $q_i \otimes q_j$ has degree $(\deg(q_i), \deg(q_j))$. The subset of homogeneous elements $\{q_i \otimes q_j, q_3 \otimes q_3 \mid i, j = 0, 1, 2\}$ spans $H_4(\mathbb{F}, *)$, and hence the 10-dimensional space $H_4(\mathbb{F}, *)$ is also \mathbb{Z}_2^4 -graded. Moreover, as the \mathbb{Z}_2 -grading on \mathcal{K} is given by $\mathcal{K}_{\bar{0}} = \langle 1 = e_1 + e_2 \rangle$ and $\mathcal{K}_{\bar{1}} = \langle e_1 - e_2 \rangle$, then $H_4(\mathcal{K}, *) \subset \mathrm{Mat}_{2 \times 2}(\mathbb{F}) \otimes \mathrm{Mat}_{2 \times 2}(\mathbb{F}) \otimes \mathcal{K}$ is spanned by the following subset of homogeneous elements for the \mathbb{Z}_2^5 -grading: $\{q_i \otimes q_j \otimes 1, q_3 \otimes q_3 \otimes 1 \mid i, j = 0, 1, 2\} \cup \{q_i \otimes q_3 \otimes (e_1 - e_2), q_3 \otimes q_i \otimes (e_1 - e_2) \mid i = 0, 1, 2\}$. The remaining cases are dealt with in the same way.

3.5 Gradings on Lie algebras obtained from Kantor's and Steinberg's constructions

Recall that we can get the exceptional Lie algebras of the E series by means of Kantor's construction applied to the structurable algebras $\mathfrak{C}\mathfrak{D}(H_4(C, *))$, for an associative Hurwitz algebra C . In turn, these structurable algebras are obtained from the Jordan algebras $J = H_4(C, *)$ by means of the Cayley-Dickson

doubling process explained in Example 2.2. This doubling process provides a \mathbb{Z}_2 -grading as usual, with even part J and odd part vJ , clearly compatible with any grading on J . At the same time, any G -grading on a structurable algebra $(A, -)$ provides a $\mathbb{Z} \times G$ -grading on $\mathfrak{Kan}(A, -)$ and a $\mathbb{Z}_2^2 \times G$ -grading on $\mathcal{U}(A, -)$ (which is isomorphic to $\mathfrak{Kan}(A, -)$). Thus we have another source of gradings on our Lie algebras. If Kantor's construction is applied to $\mathfrak{CD}(H_4(C, *))$, and the \mathbb{Z}_2 -grading induced by the Cayley-Dickson doubling process and the finite gradings on Section 3.4 are combined, we obtain gradings on:

(6g9) $\mathfrak{e}_6 = \mathfrak{Kan}(\mathfrak{CD}(H_4(\mathbb{F})))$ by the group $\mathbb{Z} \times \mathbb{Z}_2^5$, of type (73, 0, 0, 0, 1);

(7g7) $\mathfrak{e}_7 = \mathfrak{Kan}(\mathfrak{CD}(H_4(\mathcal{K})))$ by the group $\mathbb{Z} \times \mathbb{Z}_2^6$, of type (127, 0, 0, 0, 0, 1);

(8g7) $\mathfrak{e}_8 = \mathfrak{Kan}(\mathfrak{CD}(H_4(\mathcal{Q})))$ by the group $\mathbb{Z} \times \mathbb{Z}_2^7$, of type (241, 0, 0, 0, 0, 0, 1).

In the same vein, but using the infinite gradings on Section 3.4, we get gradings on:

(6g10) $\mathfrak{e}_6 = \mathfrak{Kan}(\mathfrak{CD}(H_4(\mathbb{F})))$ by the group $\mathbb{Z}^2 \times \mathbb{Z}_2^3$, of type (60, 7, 0, 1);

(7g8) $\mathfrak{e}_7 = \mathfrak{Kan}(\mathfrak{CD}(H_4(\mathcal{K})))$ by the group $\mathbb{Z}^2 \times \mathbb{Z}_2^4$, of type (102, 13, 0, 0, 1);

(8g8) $\mathfrak{e}_8 = \mathfrak{Kan}(\mathfrak{CD}(H_4(\mathcal{Q})))$ by the group $\mathbb{Z}^2 \times \mathbb{Z}_2^5$, of type (180, 31, 0, 0, 0, 1).

Moreover, if we use Steinberg's construction applied to $\mathfrak{CD}(H_4(C, *))$, the \mathbb{Z}_2 -grading induced by the Cayley-Dickson doubling process and the finite gradings in Section 3.4 can be combined to get gradings on:

(6g11) $\mathfrak{e}_6 = \mathcal{U}(\mathfrak{CD}(H_4(\mathbb{F})))$ by the group \mathbb{Z}_2^7 , of type (72, 0, 0, 0, 0, 1);

(7g9) $\mathfrak{e}_7 = \mathcal{U}(\mathfrak{CD}(H_4(\mathcal{K})))$ by the group \mathbb{Z}_2^8 , of type (126, 0, 0, 0, 0, 0, 1);

(8g9) $\mathfrak{e}_8 = \mathcal{U}(\mathfrak{CD}(H_4(\mathcal{Q})))$ by the group \mathbb{Z}_2^9 , of type (240, 0, 0, 0, 0, 0, 0, 1).

It is not difficult to see that the $\mathbb{Z} \times \mathbb{Z}_2^5$ -grading (respectively $\mathbb{Z} \times \mathbb{Z}_2^6$ and $\mathbb{Z} \times \mathbb{Z}_2^7$) obtained in $\mathfrak{e}_6 = \mathcal{U}(\mathfrak{CD}(H_4(\mathbb{F})))$ (respectively $\mathfrak{e}_7 = \mathcal{U}(\mathfrak{CD}(H_4(\mathcal{K})))$ and $\mathfrak{e}_8 = \mathcal{U}(\mathfrak{CD}(H_4(\mathcal{Q})))$) is isomorphic to the grading (6g9) (respectively (7g7) and (8g7)).

Remark 3.1. As we know [17] about the existence of a $\mathbb{Z}_4 \times \mathbb{Z}_2^4$ -grading on \mathfrak{e}_6 , of type (48, 13, 0, 1), we would like to find a \mathbb{Z}_4 -grading on the Lie algebra obtained by means of Kantor's construction attached to a structurable algebra. That can be done for $A = \mathfrak{CD}(H_4(C)) = J \oplus vJ$, with $J = H_4(C, *)$ the Jordan algebra of hermitian matrices with coefficients in an associative Hurwitz algebra

C. In such a case, $S = \mathbb{F}v$, and it is a straightforward computation, taking into account Equations (4) and (5), that $L = \mathfrak{Kan}(A)$ is \mathbb{Z}_4 -graded as follows:

$$\begin{aligned} L_{\bar{0}} &= \mathbb{F}v \oplus (V_{J,J} + V_{vJ,vJ}) \oplus \mathbb{F}v, \\ L_{\bar{1}} &= J \oplus (vJ)^\sim, \\ L_{\bar{2}} &= V_{J,vJ}, \\ L_{\bar{3}} &= J^\sim \oplus vJ. \end{aligned}$$

This grading is compatible with any grading on J , so that we can combine it with the gradings described in Section 3.4 to get gradings on $\mathfrak{e}_6 = \mathfrak{Kan}(\mathfrak{CD}(H_4(\mathbb{F})))$ by the group $\mathbb{Z}_4 \times \mathbb{Z}_2^4$, on $\mathfrak{e}_7 = \mathfrak{Kan}(\mathfrak{CD}(H_4(\mathcal{K})))$ by the group $\mathbb{Z}_4 \times \mathbb{Z}_2^5$, and on $\mathfrak{e}_8 = \mathfrak{Kan}(\mathfrak{CD}(H_4(\mathcal{Q})))$ by the group $\mathbb{Z}_4 \times \mathbb{Z}_2^6$. But these are not fine!, so we must continue the search in order to explain several gradings by groups with factors \mathbb{Z}_4 .

3.6 A $\mathbb{Z}_4 \times \mathbb{Z}_2^3$ -grading on the Jordan algebra $H_4(\mathcal{Q})$ and related gradings on the exceptional Lie algebras

A graded division (associative) algebra D is a graded algebra such that every homogeneous element is invertible. If the support of such a grading is H and G is a group containing H as a subgroup, and we have a G -graded right D -module V (that is, $V_g D_h \subset V_{g+h}$ for any $g \in G$ and $h \in H$), then the division property of D forces V to be a free right D -module containing bases consisting of homogeneous elements, according to [21, §2]. Then we have a G -grading induced on $R = \text{End}_D(V)$ given by $f \in R_g$ if $f(V_{g'}) \subset V_{g+g'}$ for any $g' \in G$.

Let $\tau: D \rightarrow D$ be a graded antiautomorphism, that is, τ is an antiautomorphism with $\tau(D_h) = D_h$ for any $h \in H$ (which implies that necessarily τ is an involution, that is $\tau^2 = \text{id}_D$). Let $b: V \times V \rightarrow D$ a sesquilinear form (b is \mathbb{F} -bilinear, $b(v_1, v_2) = \tau(b(v_2, v_1))$ and $b(v_1, v_2 d) = b(v_1, v_2) d$ for any $v_1, v_2 \in V$ and $d \in D$) compatible with the grading, that is, $b(V_g, V_{g'}) \subset D_{g+g'}$. Let $*$ be the adjoint relative to this form ($b(f(v_1), v_2) = b(v_1, f^*(v_2))$ if $f \in R$). The point is that the sets of hermitian and skew-hermitian elements $H(R, *) = \{f \in R \mid f^* = f\}$ and $K(R, *) = \{f \in R \mid f^* = -f\}$ are graded subspaces. Moreover, it is proven in [21] that essentially all the gradings in $K(R, *)$ and in $H(R, *)$ are obtained in this way.

Note that the quaternion algebra $\mathcal{Q} = \text{Mat}_{2 \times 2}(\mathbb{F})$ is a \mathbb{Z}_2^2 -graded division algebra with the grading given by the matrices in Equation (14),

$$\mathcal{Q}_{(\bar{0}, \bar{0})} = \mathbb{F}1, \quad \mathcal{Q}_{(\bar{1}, \bar{0})} = \mathbb{F}q_1, \quad \mathcal{Q}_{(\bar{0}, \bar{1})} = \mathbb{F}q_2, \quad \mathcal{Q}_{(\bar{1}, \bar{1})} = \mathbb{F}q_3. \quad (15)$$

There are two involutions compatible with this grading. The involution τ^o given by

$$q_1^{\tau^o} = q_1, \quad q_2^{\tau^o} = q_2, \quad q_3^{\tau^o} = -q_3,$$

is the usual transpose involution (an orthogonal involution); while the involution $\tau^s = -$ given by

$$\bar{q}_i = -q_i, \quad \forall i = 1, 2, 3,$$

is the standard conjugation of the quaternion algebra \mathcal{Q} (a symplectic involution).

Take $(D = \mathcal{Q}, \tau^o)$ as above but with the following grading, equivalent to (15),

$$\mathcal{Q}_{(\bar{0}, \bar{0})} = \mathbb{F}1, \quad \mathcal{Q}_{(\bar{2}, \bar{0})} = \mathbb{F}q_1, \quad \mathcal{Q}_{(\bar{0}, \bar{1})} = \mathbb{F}q_2, \quad \mathcal{Q}_{(\bar{2}, \bar{1})} = \mathbb{F}q_3.$$

Take $B = \{v_0, v_1\}$ a homogeneous D -basis in V , a graded right free D -module of dimension 2, with $\deg(v_0) = (\bar{0}, \bar{0})$ and $\deg(v_1) = (\bar{1}, \bar{0})$. We have chosen the degrees such that the sesquilinear form $b: V \times V \rightarrow D$ given by the matrix $A = \begin{pmatrix} 1 & 0 \\ 0 & q_1 \end{pmatrix}$ relative to B is compatible with the grading (since $2 \deg(v_0) = \deg(1)$ and $2 \deg(v_1) = \deg(q_1)$). Now for $x = \begin{pmatrix} p_1 & p_2 \\ p_3 & p_4 \end{pmatrix} \in \text{Mat}_{2 \times 2}(\mathcal{Q}) \simeq \text{End}_{\mathcal{Q}}(V) = R$, we have $x^* = A^{-1}(\tau^o(x))^t A = \begin{pmatrix} \tau^o(p_1) & \tau^o(p_3)q_1 \\ -q_1\tau^o(p_2) & -q_1\tau^o(p_4)q_1 \end{pmatrix}$, and hence

$$K(\text{Mat}_{2 \times 2}(\mathcal{Q}), *) = \left\{ \begin{pmatrix} \alpha q_3 & -\tau^o(p)q_1 \\ p & \beta q_2 \end{pmatrix} \mid \alpha, \beta \in \mathbb{F}, p \in \mathcal{Q} \right\} =: K$$

inherits the $\mathbb{Z}_4 \times \mathbb{Z}_2$ -grading, with 6 pieces of dimension one,

$$K = K_{(\bar{2}, \bar{1})} \oplus K_{(\bar{0}, \bar{1})} \oplus K_{(\bar{1}, \bar{0})} \oplus K_{(\bar{3}, \bar{0})} \oplus K_{(\bar{1}, \bar{1})} \oplus K_{(\bar{3}, \bar{1})},$$

and also, for $H := H(\text{Mat}_{2 \times 2}(\mathcal{Q}), *)$,

$$H = \left\{ \begin{pmatrix} p_1 & \tau^o(p)q_1 \\ p & p_2 \end{pmatrix} \mid p_1 \in \langle 1, q_1, q_2 \rangle, p_2 \in \langle 1, q_1, q_3 \rangle, p \in \mathcal{Q} \right\}$$

inherits the $\mathbb{Z}_4 \times \mathbb{Z}_2$ -grading,

$$H = H_{(\bar{2}, \bar{1})} \oplus H_{(\bar{0}, \bar{1})} \oplus H_{(\bar{1}, \bar{0})} \oplus H_{(\bar{3}, \bar{0})} \oplus H_{(\bar{1}, \bar{1})} \oplus H_{(\bar{3}, \bar{1})} \oplus H_{(\bar{0}, \bar{0})} \oplus H_{(\bar{2}, \bar{0})},$$

of type (6, 2) since all the above homogeneous components are one-dimensional except for two of them, namely,

$$H_{(\bar{0}, \bar{0})} = \left\langle \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \right\rangle, \quad H_{(\bar{2}, \bar{0})} = \left\langle \begin{pmatrix} q_1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & q_1 \end{pmatrix} \right\rangle.$$

Next we identify $\text{Mat}_{4 \times 4}(\mathcal{Q})$ with $\text{Mat}_{2 \times 2}(\mathbb{F}) \otimes \text{Mat}_{2 \times 2}(\mathbb{F}) \otimes \mathcal{Q}$ and hence with $\mathcal{Q} \otimes \text{Mat}_{2 \times 2}(\mathcal{Q}) = \mathcal{Q} \otimes R$, and consider here the (symplectic) involution

given by $\tau^s \otimes *$. The Jordan algebra $J = H_4(\mathcal{Q})$ lives here as $J = \{q \otimes x \in \mathcal{Q} \otimes R \mid \tau^s(q) \otimes x^* = q \otimes x\}$, which can be identified with $K(\mathcal{Q}, \tau^s) \otimes K(\text{Mat}_{2 \times 2}(\mathcal{Q}), *) \oplus H(\mathcal{Q}, \tau^s) \otimes H(\text{Mat}_{2 \times 2}(\mathcal{Q}), *)$. In this way, by combining the \mathbb{Z}_2^2 -grading on \mathcal{Q} given by Equation (15) with the above $\mathbb{Z}_4 \times \mathbb{Z}_2$ -grading on R , we get a $\mathbb{Z}_4 \times \mathbb{Z}_2^3$ -grading on $J = H_4(\mathcal{Q})$ of type $3(6, 0) + 1(6, 2) = (24, 2)$. Also we get a $\mathbb{Z}_4 \times \mathbb{Z}_2^3$ -grading on $K(\mathcal{Q} \otimes R, \tau^s \otimes *) \cong \text{Der}(H_4(\mathcal{Q}))$ (a Lie algebra of type \mathfrak{e}_4) of type $1(6, 0) + 3(6, 2) = (24, 6)$.

Of course this grading on J induces a $\mathbb{Z}_4 \times \mathbb{Z}_2^4$ -grading on the structurable algebra $\mathfrak{CD}(H_4(\mathcal{Q}))$, of type $(48, 4)$. Now note that, according to [3],

$$\text{Der}(\mathfrak{CD}(H_4(\mathcal{Q})), -) \simeq \mathfrak{e}_6, \quad \mathfrak{Instr}(\mathfrak{CD}(H_4(\mathcal{Q}))) \simeq \mathfrak{e}_7, \quad \mathcal{U}(\mathfrak{CD}(H_4(\mathcal{Q}))) \simeq \mathfrak{e}_8. \quad (16)$$

In particular, every G -grading on $\mathfrak{CD}(H_4(\mathcal{Q}))$ induces a G -grading on \mathfrak{e}_6 , a $G \times \mathbb{Z}_2$ -grading on \mathfrak{e}_7 and a $G \times \mathbb{Z}_2^2$ -grading on \mathfrak{e}_8 . In our case we get gradings on:

(6g12) $\mathfrak{e}_6 = \text{Der}(\mathfrak{CD}(H_4(\mathcal{Q})))$ by the group $\mathbb{Z}_4 \times \mathbb{Z}_2^4$, of type $(48, 13, 0, 1)$;

(7g10) $\mathfrak{e}_7 = \mathfrak{Instr}(\mathfrak{CD}(H_4(\mathcal{Q})))$ by the group $\mathbb{Z}_4 \times \mathbb{Z}_2^5$, of type $(98, 15, 0, 0, 1)$;

(8g10) $\mathfrak{e}_8 = \mathcal{U}(\mathfrak{CD}(H_4(\mathcal{Q})))$ by the group $\mathbb{Z}_4 \times \mathbb{Z}_2^6$, of type $(192, 25, 0, 0, 0, 1)$.

3.7 A \mathbb{Z}_4^3 -grading on the structurable algebra $\mathfrak{CD}(H_4(\mathcal{Q}))$

There is a \mathbb{Z}_4^3 -grading on the structurable algebra $\mathcal{A} = \mathfrak{CD}(H_4(\mathcal{Q}))$ which is not explained in terms of the Cayley-Dickson process. This is a very interesting grading in which every nonzero homogeneous component is one-dimensional. Let us describe it. The information is extracted from [8].

Identify, as in Equation (13), $\text{Mat}_{4 \times 4}(\mathcal{Q})$ with $\text{Mat}_{4 \times 4}(\mathbb{F}) \otimes \mathcal{Q}$. The involution $(q_{ij})^* = (\bar{q}_{ji})$ in $\text{Mat}_{4 \times 4}(\mathcal{Q})$ is, under such correspondence, the tensor product of the matrix transpose on $\text{Mat}_{4 \times 4}(\mathbb{F})$ and the standard involution on \mathcal{Q} (under the identification of \mathcal{Q} with $\text{Mat}_{2 \times 2}(\mathbb{F})$, this involution acts as follows: $\bar{E}_{11} = E_{22}$, $\bar{E}_{22} = E_{11}$, $\bar{E}_{12} = -E_{12}$, $\bar{E}_{21} = -E_{21}$). In particular, the Jordan subalgebra of symmetric elements $H_4(\mathcal{Q}) = \{q = (q_{ij}) \in \text{Mat}_{4 \times 4}(\mathcal{Q}) \mid q^* = q\}$ is identified with $H_4(\mathbb{F}) \otimes \langle E_{11} + E_{22} \rangle \oplus K_4(\mathbb{F}) \otimes \langle E_{11} - E_{22}, E_{12}, E_{21} \rangle$, where $H_4(\mathbb{F})$ and $K_4(\mathbb{F})$ denote the subspaces of symmetric and skewsymmetric matrices of size 4, respectively, and hence with

$$\mathcal{J} = \left\{ \left(\begin{array}{cc} z & x \\ y & z^t \end{array} \right) \mid x = -x^t, y = -y^t, x, y, z \in \text{Mat}_{4 \times 4}(\mathbb{F}) \right\}.$$

Consider the following \mathbb{Z}_4 -grading on the structurable algebra $\mathcal{A} = \mathcal{J} \oplus v\mathcal{J}$:

$$\begin{aligned} \mathcal{A}_{\bar{0}} &= \left\{ \begin{pmatrix} z & 0 \\ 0 & z^t \end{pmatrix} \mid z \in \text{Mat}_{4 \times 4}(\mathbb{F}) \right\}, \\ \mathcal{A}_{\bar{1}} &= \left\{ \begin{pmatrix} 0 & x \\ 0 & 0 \end{pmatrix} + v \begin{pmatrix} 0 & 0 \\ y & 0 \end{pmatrix} \mid x, y \in K_4(\mathbb{F}) \right\}, \\ \mathcal{A}_{\bar{2}} &= v\mathcal{A}_{\bar{0}}, \\ \mathcal{A}_{\bar{3}} &= \left\{ v \begin{pmatrix} 0 & x \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ y & 0 \end{pmatrix} \mid x, y \in K_4(\mathbb{F}) \right\}. \end{aligned} \quad (17)$$

Note that if $u \in \text{GL}(4, \mathbb{F})$ is an invertible matrix, we can consider the following automorphism of $H_4(\mathcal{Q})$,

$$\Psi(u) : \begin{pmatrix} z & x \\ y & z^t \end{pmatrix} \mapsto \begin{pmatrix} uzu^{-1} & uxu^t \\ (u^{-1})^t y u^{-1} & (uzu^{-1})^t \end{pmatrix},$$

which extends to \mathcal{A} in a natural way (also denoted $\Psi(u)$). For any natural number n , let ξ be a primitive n th root of 1 and consider the following matrices

$$P_n = \begin{pmatrix} 1 & 0 & \dots & \dots & 0 \\ 0 & \xi & 0 & \dots & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & \dots & 0 & \xi^{n-2} & 0 \\ 0 & \dots & \dots & 0 & \xi^{n-1} \end{pmatrix}, \quad Q_n = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & \dots & 0 & 0 & 1 \\ 1 & 0 & \dots & \dots & 0 \end{pmatrix}, \quad (18)$$

also called Pauli matrices (P_n sometimes will be denoted by P_ξ). Take now the Pauli matrices $X = P_4$ and $Y = Q_4$ and note that then $\Psi(X)$ and $\Psi(Y)$ are order four automorphisms of \mathcal{A} (which neither commute nor anticommute).

Also, consider for any skew-symmetric 4×4 matrix, its Pfaffian adjoint \hat{x} :

$$x = \begin{pmatrix} 0 & \alpha & \beta & \gamma \\ -\alpha & 0 & \delta & \varepsilon \\ -\beta & -\delta & 0 & \zeta \\ -\gamma & -\varepsilon & -\zeta & 0 \end{pmatrix} \in K_4(\mathbb{F}), \quad \hat{x} = \begin{pmatrix} 0 & -\zeta & \varepsilon & -\delta \\ \zeta & 0 & -\gamma & \beta \\ -\varepsilon & \gamma & 0 & -\alpha \\ \delta & -\beta & \alpha & 0 \end{pmatrix}.$$

(Note that this differs from [8], where $-\hat{x}$ is considered.) Now consider the order 4 automorphism $\pi: \mathcal{A} \rightarrow \mathcal{A}$, whose restriction to $\mathcal{A}_{\bar{0}} \oplus \mathcal{A}_{\bar{2}}$ is the identity, and such that:

$$\pi \begin{pmatrix} 0 & x \\ y & 0 \end{pmatrix} = v \begin{pmatrix} 0 & -\hat{y} \\ \hat{x} & 0 \end{pmatrix}, \quad \pi \left(v \begin{pmatrix} 0 & x \\ y & 0 \end{pmatrix} \right) = \begin{pmatrix} 0 & -\hat{y} \\ \hat{x} & 0 \end{pmatrix},$$

on $\mathcal{A}_{\bar{1}} \oplus \mathcal{A}_{\bar{3}}$. If $\xi \in \mathbb{F}$ is chosen such that $\xi^2 = \mathbf{i}$, then $\pi\Psi(\xi X)$ and $\Psi(Y)$ are two order 4 commuting automorphisms that preserve the \mathbb{Z}_4 -grading given by Equation (17), and a \mathbb{Z}_4^3 -grading on \mathcal{A} is obtained whose homogeneous components

are the intersection of the homogeneous components of the \mathbb{Z}_4 -grading with the common eigenspaces for $\pi\Psi(\xi X)$ and $\Psi(Y)$.

Remark 3.2. This structurable algebra of dimension 56 (the only simple one of such dimension) has a model which is better known. It is defined on the vector space

$$\begin{pmatrix} \mathbb{F} & \mathbb{A} \\ \mathbb{A} & \mathbb{F} \end{pmatrix}$$

with multiplication given by

$$\begin{pmatrix} \alpha_1 & x_1 \\ x'_1 & \beta_1 \end{pmatrix} \begin{pmatrix} \alpha_2 & x_2 \\ x'_2 & \beta_2 \end{pmatrix} = \begin{pmatrix} \alpha_1\alpha_2 + T(x_1, x'_2) & \alpha_1x_2 + \beta_2x_1 + x'_1 \times x'_2 \\ \alpha_2x'_1 + \beta_1x'_2 + x_1 \times x_2 & \beta_1\beta_2 + T(x_2, x'_1) \end{pmatrix},$$

where T denotes the map $T: \mathbb{A} \times \mathbb{A} \rightarrow \mathbb{F}$ given by $T(x, y) = T(xy)$ and \times denotes the so called Freudenthal cross product defined by $T(x \times y, z) = N(x, y, z)$ if $x, y, z \in \mathbb{A}$. The involution is given by

$$\overline{\begin{pmatrix} \alpha & x \\ x' & \beta \end{pmatrix}} = \begin{pmatrix} \beta & x \\ x' & \alpha \end{pmatrix}.$$

Although this is isomorphic (as an algebra with involution), to $\mathfrak{CD}(H_4(\mathcal{Q}))$, it was previously studied as an example of Brown algebra. Garibaldi [30] discusses the connections between this algebra and the groups of types E_6 and E_7 .

3.8 More gradings on the exceptional Lie algebras by groups with factors \mathbb{Z}_4

The \mathbb{Z}_4^3 -grading above on $\mathfrak{CD}(H_4(\mathcal{Q}))$ immediately induces the following gradings on:

(8g11) $\mathfrak{e}_8 = \mathfrak{kan}(\mathfrak{CD}(H_4(\mathcal{Q})))$ by the group $\mathbb{Z} \times \mathbb{Z}_4^3$, of type (123, 40, 15);

(8g12) $\mathfrak{e}_8 = \mathcal{U}(\mathfrak{CD}(H_4(\mathcal{Q})))$ by the group $\mathbb{Z}_2^2 \times \mathbb{Z}_4^3$, of type (216, 14, 0, 1);

if we take into consideration the \mathbb{Z} -grading provided by Kantor's construction and the \mathbb{Z}_2^2 -grading provided by Steinberg's construction.

Furthermore, recall that the algebras \mathfrak{e}_6 and \mathfrak{e}_7 can be obtained too from the structurable algebra $\mathfrak{CD}(H_4(\mathcal{Q}))$ as in (16). Hence we get also gradings on:

(6g13) $\mathfrak{e}_6 = \text{Der}(\mathfrak{CD}(H_4(\mathcal{Q})))$ by the group \mathbb{Z}_4^3 , of type (48, 15);

(7g11) $\mathfrak{e}_7 = \mathfrak{Instr}(\mathfrak{CD}(H_4(\mathcal{Q})))$ by the group $\mathbb{Z}_4^3 \times \mathbb{Z}_2$, of type (102, 14, 1).

The restriction of the \mathbb{Z}_4^3 -grading on $\mathfrak{CD}(H_4(\mathcal{Q}))$ to its structurable subalgebra $\mathfrak{CD}(H_4(\mathcal{K}))$ provides a $\mathbb{Z}_4^2 \times \mathbb{Z}_2$ -grading on $\mathfrak{CD}(H_4(\mathcal{K}))$, which of course can be used to get gradings on:

(7g12) $\mathfrak{e}_7 = \mathfrak{kan}(\mathfrak{CD}(H_4(\mathcal{K})))$ by the group $\mathbb{Z} \times \mathbb{Z}_4^2 \times \mathbb{Z}_2$, of type (67, 27, 4);

(7g13) $\mathfrak{e}_7 = \mathcal{U}(\mathfrak{CD}(H_4(\mathcal{K})))$ by the group $\mathbb{Z}_2^2 \times \mathbb{Z}_4^2 \times \mathbb{Z}_2 = \mathbb{Z}_2^3 \times \mathbb{Z}_4^2$, of type (123, 3, 0, 1).

3.9 A fine \mathbb{Z}_5^3 -grading

There is a \mathbb{Z}_5^3 -grading on \mathfrak{e}_8 which seems not to be related with any of the previous constructions or structures. This grading appears in several contexts (for instance, [1] and [20], and lately in [42]), due to its interesting properties: the zero homogeneous component is trivial (as in any fine grading by a finite group, see [16, Corollary 5]) and all the other homogeneous components (in this case 124) have the same dimension (so that in this case such dimension must be 2) and consist of semisimple elements ([17, Lemma 1]). Moreover, given any $0 \neq g \in \mathbb{Z}_5^3$, the subspace $\bigoplus_{i=1}^4 (\mathfrak{e}_8)_{ig}$ is a Cartan subalgebra. The following description can be found in [20].

Let V_1 and V_2 be two vector spaces over \mathbb{F} of dimension 5 and let us consider the following \mathbb{Z}_5 -graded vector space $L = L_{\bar{0}} \oplus L_{\bar{1}} \oplus L_{\bar{2}} \oplus L_{\bar{3}} \oplus L_{\bar{4}}$, for

$$\begin{aligned} L_{\bar{0}} &= \mathfrak{sl}(V_1) \oplus \mathfrak{sl}(V_2), \\ L_{\bar{1}} &= V_1 \otimes \wedge^2 V_2, \\ L_{\bar{2}} &= \wedge^2 V_1 \otimes \wedge^4 V_2, \\ L_{\bar{3}} &= \wedge^3 V_1 \otimes V_2, \\ L_{\bar{4}} &= \wedge^4 V_1 \otimes \wedge^3 V_2. \end{aligned} \tag{19}$$

We can endow L with a structure of \mathbb{Z}_5 -graded Lie algebra, with the natural action of the semisimple algebra $L_{\bar{0}}$ on each of the other homogeneous components. The brackets involving elements in different homogeneous components are given by suitable scalar multiples of the only $L_{\bar{0}}$ -invariant maps from $L_{\bar{i}} \times L_{\bar{j}} \rightarrow L_{\bar{i}+\bar{j}}$ (these scalars have been computed explicitly in [13]). The Lie algebra defined in this way is simple of dimension 248, and hence it provides a linear model of \mathfrak{e}_8 . The philosophy of this kind of linear models can be found in [46, Chapter 5, §2].

Let $\xi \in \mathbb{F}$ be a primitive fifth root of 1, and take B_1 and B_2 bases of V_1 and V_2 respectively, and endomorphisms $b_1, c_1 \in \text{End}(V_1)$ and $b_2, c_2 \in \text{End}(V_2)$ whose coordinate matrices in the bases B_i are

$$\begin{aligned} b_1 &= P_\xi, & c_1 &= Q_5, \\ b_2 &= P_{\xi^2}, & c_2 &= Q_5, \end{aligned}$$

defined as in Equation (18).

Now the unique automorphisms $\Psi, \Psi' \in \text{Aut}(L)$ whose restrictions to $L_{\bar{1}}$ are given by

$$\begin{aligned}\Psi|_{L_{\bar{1}}} &= b_1 \otimes \wedge^2 b_2, \\ \Psi'|_{L_{\bar{1}}} &= c_1 \otimes \wedge^2 c_2,\end{aligned}$$

are order 5 automorphisms which commute with the automorphism producing the \mathbb{Z}_5 -grading on L given by Equation (19). Thus we obtain the desired grading by the group \mathbb{Z}_5^3 , of type $(0, 124)$.

3.10 Gradings induced from other linear models

We would like to explain a little bit the history of the search for the gradings described in Section 3.8, which eventually lead to the quest for the grading by \mathbb{Z}_4^3 on the simple structurable algebra of dimension 56.

Consider the chain $E_6 \subset E_7 \subset E_8$ of exceptional groups. The maximal abelian subgroup \mathbb{Z}_4^3 of E_6 is then also an abelian subgroup of E_7 and also of E_8 , predictably non-toral. This forced us to consider the order 4 automorphisms of \mathfrak{e}_7 and \mathfrak{e}_8 . First note that if we look at the subgroup \mathbb{Z}_4^3 of E_6 , the three copies of \mathbb{Z}_4 involved do not play the same role. One comes from ϑ , an outer automorphism of \mathfrak{e}_6 producing the grading


$$\mathfrak{e}_6 = (\mathfrak{a}_3 \oplus \mathfrak{sl}(V)) \oplus (V(2\lambda_1) \otimes V) \oplus (V(2\lambda_2) \otimes \mathbb{F}) \oplus (V(2\lambda_3) \otimes V), \quad (20)$$

for a two-dimensional vector space V , where the λ_i 's are the fundamental dominant weights for \mathfrak{a}_3 . The other copies of \mathbb{Z}_4 , restricted to \mathfrak{a}_3 , produce the \mathbb{Z}_4^2 -grading obtained by means of Pauli matrices. More precisely, they correspond to the group

$$\langle (P_4, P_2), (Q_4, Q_2) \rangle \subset \frac{\text{SL}(4) \times \text{SL}(2)}{\langle (iI_4, -I_2) \rangle} \simeq \text{Cent}_{E_6} \langle \vartheta \rangle.$$

The \mathbb{Z}_4^3 -grading on \mathfrak{e}_6 is easily handled in this way, since we obtain concrete descriptions of the homogeneous components in terms of tensors of the natural representations of $\mathfrak{sl}(4)$ and $\mathfrak{sl}(2)$.

Inspired by this, one can consider the automorphism of \mathfrak{e}_7 obtained by removing the black node of the extended Dynkin diagram (see [39, Chapter 8])




which produces a \mathbb{Z}_4 -grading $\mathcal{L} = \mathcal{L}_0 \oplus \mathcal{L}_1 \oplus \mathcal{L}_2 \oplus \mathcal{L}_3$ on the Lie algebra $\mathcal{L} \cong \mathfrak{e}_7$, where

$$\begin{aligned}\mathcal{L}_0 &= \mathfrak{sl}(W_1) \oplus \mathfrak{sl}(V) \oplus \mathfrak{sl}(W_2), \\ \mathcal{L}_1 &= W_1 \otimes V \otimes W_2, \\ \mathcal{L}_2 &= \bigwedge^2 W_1 \otimes \mathbb{F} \otimes \bigwedge^2 W_2, \\ \mathcal{L}_3 &= \bigwedge^3 W_1 \otimes V \otimes \bigwedge^3 W_2,\end{aligned}$$

for $W_1 = W_2$ and V vector spaces of dimensions 4 and 2 respectively. (This gives the structure of the homogeneous components as modules for \mathcal{L}_0 .) We consider now the order four automorphisms whose restrictions to \mathcal{L}_1 are $P_4 \otimes P_2 \otimes P_4$, $Q_4 \otimes Q_2 \otimes Q_4$, with P_n and Q_n defined as in Equation (18), and the order 2 automorphism determined by $w_1 \otimes v \otimes w_2 \mapsto w_2 \otimes v \otimes w_1$. In this way, a $\mathbb{Z}_4^3 \times \mathbb{Z}_2$ -grading on \mathfrak{e}_7 is obtained (equivalent to **(7g11)**).

The same kind of arguments can be used to study \mathbb{Z}_4 -gradings on \mathfrak{e}_8 . Again, remove the black node of the extended Dynkin diagram of \mathfrak{e}_8



to get an automorphism $\vartheta \in \text{Aut}(\mathfrak{e}_8)$ producing a \mathbb{Z}_4 -grading $\mathcal{L} = \mathcal{L}_0 \oplus \mathcal{L}_1 \oplus \mathcal{L}_2 \oplus \mathcal{L}_3$ on the Lie algebra $\mathcal{L} \cong \mathfrak{e}_8$ where, as \mathcal{L}_0 -modules, we have:

$$\begin{aligned}\mathcal{L}_0 &= \mathfrak{sl}(W) \oplus \mathfrak{sl}(V), \\ \mathcal{L}_1 &= \bigwedge^2 W \otimes V, \\ \mathcal{L}_2 &= \bigwedge^4 W \otimes \mathbb{F}, \\ \mathcal{L}_3 &= \bigwedge^6 W \otimes V,\end{aligned}$$

for W and V vector spaces of dimensions 8 and 2 respectively. The centralizer can be checked to be

$$\text{Cent}_{E_8}(\vartheta) \simeq \frac{\text{PSL}(8) \times \text{PSL}(2)}{\langle (iI_8, -I_2) \rangle},$$

where the automorphism ϑ corresponds to the class $[(\xi I_8, I_2)]$ for $\xi \in \mathbb{F}$ such that $\xi^2 = i$. The MAD-groups of E_8 containing ϑ are MAD-groups of $\text{Cent}_{E_8}(\vartheta)$. In particular, if we take ϑ together with

$$\begin{aligned}[(I_2 \otimes P_4, P_2)], & \quad [(P_2 \otimes I_4, I_2)], \\ [(I_2 \otimes Q_4, Q_2)], & \quad [(Q_2 \otimes I_4, I_2)],\end{aligned}$$

we obtain a $\mathbb{Z}_4^3 \times \mathbb{Z}_2^2$ -grading on \mathfrak{e}_8 (equivalent to **(8g12)**); and if we take it together with

$$[(I_2 \otimes P_4, P_2)], \quad [(I_2 \otimes Q_4, Q_2)], \quad \left[\left(\begin{pmatrix} \alpha & 0 \\ 0 & \frac{1}{\alpha} \end{pmatrix} \otimes I_4, I_2 \right) \right],$$

we obtain the product of a one-dimensional torus and \mathbb{Z}_4^3 , and hence a $\mathbb{Z}_4^3 \times \mathbb{Z}$ -grading on \mathfrak{e}_8 (equivalent to **(8gi11)**).

According to [22], this latter grading has to be induced by a fine grading with universal group \mathbb{Z}_4^3 on the simple structurable algebra of dimension 56. Unfortunately, the gradings on the structurable algebras are not yet classified, but in any case it was worth to find such a \mathbb{Z}_4^3 -grading, because it lies behind several gradings on the simple Lie algebras \mathfrak{e}_6 , \mathfrak{e}_7 and \mathfrak{e}_8 (those in Section 3.8). This was the starting point of [8].

However, the models based on the \mathbb{Z}_4^3 -grading on the simple 56-dimensional structurable algebra have a disadvantage over the linear models above, as the description of the homogeneous components involving pieces of $\mathbf{trip}(\mathcal{C}\mathcal{D}(H_4(C)))$ is not an easy task, and the type of the grading or the conjugacy classes of the automorphisms are neither easy to compute.

3.11 Conclusion

A large list of gradings on exceptional Lie algebras has been compiled here. All of them are fine and are described by their universal grading groups. We summarize them in the next result.

Theorem 3.3. *The following gradings on the simple Lie algebras of type E are all fine:*

- *The gradings on \mathfrak{e}_6 described as **(6gi)**, $i = 1, \dots, 13$, whose universal groups are: \mathbb{Z}_2^6 , \mathbb{Z}_3^4 , $\mathbb{Z}_2^3 \times \mathbb{Z}_3^2$, $\mathbb{Z}_2 \times \mathbb{Z}_3^3$, $\mathbb{Z}^2 \times \mathbb{Z}_3^2$, $\mathbb{Z} \times \mathbb{Z}_2^4$, $\mathbb{Z}^4 \times \mathbb{Z}_2$, $\mathbb{Z}^2 \times \mathbb{Z}_2^3$, $\mathbb{Z} \times \mathbb{Z}_2^5$, $\mathbb{Z}^2 \times \mathbb{Z}_2^3$, \mathbb{Z}_2^7 , $\mathbb{Z}_4 \times \mathbb{Z}_2^4$, \mathbb{Z}_4^3 .*
- *The gradings on \mathfrak{e}_7 described as **(7gi)**, $i = 1, \dots, 13$, whose universal groups are: \mathbb{Z}_2^7 , $\mathbb{Z}_2^2 \times \mathbb{Z}_3^3$, $\mathbb{Z} \times \mathbb{Z}_3^3$, $\mathbb{Z} \times \mathbb{Z}_2^5$, $\mathbb{Z}^4 \times \mathbb{Z}_2^2$, $\mathbb{Z}^3 \times \mathbb{Z}_2^3$, $\mathbb{Z} \times \mathbb{Z}_2^6$, $\mathbb{Z}^2 \times \mathbb{Z}_2^4$, \mathbb{Z}_2^8 , $\mathbb{Z}_4 \times \mathbb{Z}_2^5$, $\mathbb{Z}_4^3 \times \mathbb{Z}_2$, $\mathbb{Z} \times \mathbb{Z}_4^2 \times \mathbb{Z}_2$, $\mathbb{Z}_2^3 \times \mathbb{Z}_4^2$.*
- *The gradings on \mathfrak{e}_8 described as **(8gi)**, $i = 1, \dots, 13$, whose universal groups are: \mathbb{Z}_2^8 , \mathbb{Z}_3^5 , \mathbb{Z}_6^3 , $\mathbb{Z}^2 \times \mathbb{Z}_3^3$, $\mathbb{Z} \times \mathbb{Z}_2^6$, $\mathbb{Z}^4 \times \mathbb{Z}_2^3$, $\mathbb{Z} \times \mathbb{Z}_2^7$, $\mathbb{Z}^2 \times \mathbb{Z}_2^5$, \mathbb{Z}_2^9 , $\mathbb{Z}_4 \times \mathbb{Z}_2^6$, $\mathbb{Z} \times \mathbb{Z}_4^3$, $\mathbb{Z}_2^2 \times \mathbb{Z}_4^3$, \mathbb{Z}_5^3 .*

Conjecture: We think that these gradings exhaust the list of fine gradings, up to equivalence, on \mathfrak{e}_6 , \mathfrak{e}_7 and \mathfrak{e}_8 , with the exception of the root space decompositions relative to a Cartan subalgebra.

If this conjecture were true, then there would be, up to equivalence, 14 fine gradings on each of the simple Lie algebras of type E . Therefore, there would be exactly 14 conjugacy classes of maximal abelian diagonalizable subgroups of the algebraic group $\text{Aut}(\mathfrak{e}_r)$, $r = 6, 7, 8$.

This conjecture has been proved for \mathfrak{e}_6 in [17]. The cases of \mathfrak{e}_7 and \mathfrak{e}_8 remain open. An strategy for the finite case is that every automorphism belonging to a MAD-subgroup of a connected and simply-connected group (like E_8) fixes a semisimple subalgebra, so that it corresponds to removing only one node in the extended Dynkin diagram. That fact implies that only a handful of automorphisms are possible and we are working in each case (see [14]) by studying the corresponding centralizers as in Section 3.10. The group $\text{Aut}(\mathfrak{e}_7)$ is not simply connected, but once one gets all the MAD-subgroups of E_8 , much of the work is already done. In order to deal with infinite MAD-groups, note that every grading by an infinite group is related to a grading by a root system, as proved in [22]. As these root-gradings are well known, the problem reduces to study some special gradings on the coordinate algebras. In many cases these are structurable algebras or related to them. Hence the problem is reduced to study gradings in algebras of relative low dimension (compared to the dimension of the exceptional simple Lie algebras).

The recent results by Y. Ju (*Maximal abelian subgroups of compact simple Lie groups of type E* , arXiv:1403.2680) and of A. Elduque (*Gradings on algebras over algebraically closed fields*, arXiv:1407.0480) imply that the gradings in Theorem 3.3 exhaust the fine gradings, up to equivalence, in \mathfrak{e}_6 , \mathfrak{e}_7 and \mathfrak{e}_8 .

References

- [1] A.V. ALEKSEEVSKII: *Jordan finite commutative subgroups of simple complex Lie groups*, Functional Anal. Appl. **8** (1974), no. 4, 277–279.
- [2] B.N. ALLISON: *A class of nonassociative algebras with involution containing the class of Jordan algebras*, Math. Ann. **237** (1978), 133–156.
- [3] B.N. ALLISON: *Models of isotropic simple Lie algebras*, Comm. Algebra **7** (1979), no. 17, 1835–1875.
- [4] B. ALLISON, G. BENKART AND Y. GAO: *Lie Algebras Graded by the Root Systems BC_r , $r \geq 2$* , Memoirs Amer. Math. Soc. **158** (2002), no. 751.
- [5] B.N. ALLISON AND J.R. FAULKNER: *A Cayley-Dickson Process for a class of structurable algebras*, Trans. Amer. Math. Soc. **283** (1984), no. 1, 185–210.
- [6] B.N. ALLISON AND J.R. FAULKNER: *Nonassociative Coefficient Algebras for Steinberg Unitary Lie Algebras*, J. Algebra **161** (1993), no. 1, 1–19.
- [7] D. ARANDA-ORNA, C. DRAPER AND V. GUIDO: *Weyl groups of some fine gradings on \mathfrak{e}_6* , arXiv:1308.0631.
- [8] D. ARANDA-ORNA, A. ELDUQUE AND M. KOTCHETOV: *A \mathbb{Z}_4^3 -grading on a 56-dimensional simple structurable algebra and related fine gradings on the simple Lie algebras of type E* , Comment. Math. Univ. Carolin. *55* (2014), no. 3, 285–313.
- [9] Y. BAHTURIN AND M. KOTCHETOV: *Classification of group gradings on simple Lie algebras of types A , B , C and D* , J. Algebra **324** (2010), no. 11, 2971–2989.

- [10] C.H. BARTON AND A. SUDBERY: *Magic squares and matrix models of Lie algebras*, Adv. Math. **180** (2003), no. 2, 596–647.
- [11] E. CARTAN: *Les groupes réels simples finis et continus*, Ann. Éc. Norm. **31** (1914), 263–355.
- [12] C. CHEVALLEY AND R.D. SCHAFER: *The exceptional simple Lie algebras F_4 and E_6* , Proc. Nat. Acad. Sci. **36** (1950), 137–141.
- [13] C. DRAPER: *Models of \mathfrak{e}_8* , Communication presented at the International Mediterranean Congress of Mathematics, Almería (Spain) (2005).
- [14] C. DRAPER AND A. ELDUQUE: *Maximal finite abelian subgroups of E_8* , arXiv:1312.4326.
- [15] C. DRAPER AND C. MARTÍN: *Gradings on \mathfrak{g}_2* , Linear Algebra Appl. **418** (2006), no. 1, 85–111.
- [16] C. DRAPER AND C. MARTÍN: *Gradings on the Albert algebra and on \mathfrak{f}_4* , Rev. Mat. Iberoam. **25** (2009), no. 3, 841–908.
- [17] C. DRAPER AND A. VIRUEL: *Fine gradings on \mathfrak{e}_6* , to appear in Publ. Mat., arXiv:1207.6690.
- [18] A. ELDUQUE: *The Magic Square and Symmetric Compositions*, Rev. Mat. Iberoamericana **20** (2004), 475–491.
- [19] A. ELDUQUE: *Gradings on symmetric composition algebras*, J. Algebra **322** (2009), no. 10, 3542–3579.
- [20] A. ELDUQUE: *Jordan gradings on exceptional simple Lie algebras*, Proc. Amer. Math. Soc. **137** (2009), no. 12, 4007–4017.
- [21] A. ELDUQUE: *Fine gradings on simple classical Lie algebras*, J. Algebra **324** (2010), no. 12, 3532–3571.
- [22] A. ELDUQUE: *Fine gradings and gradings by root systems on simple Lie algebras*, to appear in Rev. Mat. Iberoam.; arXiv:1303.0651.
- [23] A. ELDUQUE AND M. KOCHETOV: *Gradings on simple Lie algebras*. Mathematical Surveys and Monographs, 189. American Mathematical Society, Providence, RI; Atlantic Association for Research in the Mathematical Sciences (AARMS), Halifax, NS, 2013.
- [24] A. ELDUQUE AND H.C. MYUNG: *On flexible composition algebras*, Comm. Algebra **21** (1993), no. 7, 2481–2505.
- [25] F. ENGEL: *Ein neues, dem linearen Komplexe analoges Gebilde*, Leipz. Ber. **52** (1900), 63–76, 220–239.
- [26] H. FREUDENTHAL: *Oktaven, Ausnahmegruppen und Oktavengeometrie*, Utrecht: Mathematisch Instituut der Rijksuniversiteit **52** (1951).
- [27] H. FREUDENTHAL: *Sur le groupe exceptionnel E_7* , Nederl. Akad. Wetensch. Pro. Sr. A56 **1** (1953), 81–89.
- [28] H. FREUDENTHAL: *Lie groups in the foundations of geometry*, Advances in Math. **1** (1964), no. 2, 145–190.
- [29] W. FULTON AND J. HARRIS: *Representation Theory. A first course*. Springer Verlag New York, Inc. (1991).
- [30] R. S. GARIBALDI: *Structurable algebras and Groups of type E_6 and E_7* , Journal of Algebra **236** (2001), 651–691.

- [31] R.L. GRIESS: *Elementary abelian p -subgroups of algebraic groups*, Geom. Dedicata **39** (1991), no. 3, 253–305.
- [32] G. HANG AND D. VOGAN: *Finite Maximal Tori*, preprint (<http://www-math.mit.edu/~dav/finitetori.pdf>).
- [33] M. HAVLÍČEK, J. PATERA AND E. PELANTOVÁ: *On Lie gradings II*, Linear Algebra Appl. **277** (1998), no. 1-3, 97–125.
- [34] N. JACOBSON: *Cayley numbers and simple Lie algebras of type G* , Duke Math. J. **5** (1939), 775–783.
- [35] N. JACOBSON: *Some groups of transformations defined by Jordan algebras I*, J. Reine Angew Math. **201** (1959), 178–195.
- [36] N. JACOBSON: *Structure and representations of Jordan algebras*. American Mathematical Society Colloquium Publications, vol XXXIX. American Mathematical Society, Providence, 1964.
- [37] N. JACOBSON: *Exceptional Lie algebras*. Lecture notes in pure and applied mathematics. Marcel Dekker, Inc., New York, 1971.
- [38] V.G. KAC: *Automorphisms of finite order of semisimple Lie algebras*, Funct. Anal. Appl. **3** (1970), 252–254. (English transl.)
- [39] V.G. KAC: *Infinite-dimensional Lie algebras*, Third edition. Cambridge University Press, Cambridge, 1990.
- [40] I.L. KANTOR: *Models of Exceptional Lie algebras*, Soviet Math. Dokl. **14** (1973), no. 1, 254–258.
- [41] M. KOECHER: *Imbedding of Jordan algebras in Lie algebras, I-II*, Amer. J. Math. **89** (1967), 787–816 and **90** (1968), 476–510.
- [42] B. KOSTANT: *On some of the mathematics in Garrett Lisi's $E(8)$ Theory of Everything*, communication presented at the University of California Lie Theory Workshop, February 2008.
- [43] J.M. LANDSBERG AND L. MANIVEL: *Triality, exceptional Lie algebras and Deligne dimension formulas*, Adv. Math. **171** (2002), no. 1, 59–85.
- [44] A.G. LISI: *An Exceptionally Simple Theory of Everything*, arXiv:0711.0770.
- [45] S. OKUBO: *Pseudo-quaternion and pseudo-octonion algebras*, Hadronic J. **1** (1978), no. 4, 1250–1278.
- [46] A.L. ONISHCHNIK AND È.B. VINBERG (EDS.): *Lie Groups and Lie Algebras III*, Encyclopaedia of Mathematical Sciences **41**. Springer-Verlag, Berlin, 1991.
- [47] J. PATERA AND H. ZASSENHAUS: *On Lie gradings. I*, Linear Algebra Appl. **112** (1989), 87–159.
- [48] R.D. SCHAFER: *An introduction to nonassociative algebras*, Dover Publications Inc., New York, 1995. (Reprint of the 1966 original).
- [49] J. TITS: *Algèbres alternatives, algèbres de Jordan et algèbres de Lie exceptionnelles. I. Construction*, Indag. Math. **28** (1966), 223–237.
- [50] J. TITS: *Le plan projectif des octaves et les groupes de Lie exceptionnels*, Acad. Roy. Belg. Bull. **39** (1953), 309–329.
- [51] M.L. TOMBER: *Lie algebras of type F* , Proc. Amer. Math. Soc. **4** (1953), 759–768.
- [52] J. YU: *Maximal abelian subgroups of compact simple Lie groups*, arXiv:1211.1334.

- [53] K.A. ZHEVLAKOV, A.M. SLIN'KO, I.P. SHESTAKOV AND A.I. SHIRSHOV: Rings that are nearly associative. Pure and Applied Mathematics, vol. 104. Academic Press Inc. New York, 1982.