

General Index

PROGRAM	9
ABSTRACT BOOK	19
INVITED LECTURE	21
A focus on Italy to disentangle the earliest migration of <i>Homo sapiens</i> in Southern Europe <i>by Benazzi S.</i>	22
PRESENTATIONS	25
Climate variability in the Monte San Nicola type-section (Sicily): high-resolution calcareous plankton fluctuations over the Pliocene-Pleistocene transition <i>by Addante M.</i> , Girone A., Maiorano P., Marino M., Scopelliti G., Trotta S. & Caruso A.	26
The high dinosaur biodiversity in the Apulia carbonate platform: evidence from the Lama Balice tracksite (upper Albian, Southern Italy) <i>by Antonelli M.</i> , Petti F.M., De Sario F., Sacco E., La Perna R., Marino M., Marsico A., Petruzzelli M., Sabato L., Spalluto L. & Tropeano M.	27
Coccolith input into the reconstruction of sea surface dynamic: Iberian margin and North Atlantic Ocean <i>by Argenio C.</i> , Flores J.A., Fuertes M.A., Balestra B., Cerulo L., Cachão M. & Amore F.O.	28
Reappraisal of the upper Olduvai Beds (Tanzania): insights from a new faunal assemblage from Geolocality 83 <i>by Azzarà B.</i> , Cherin M., Boschian G., Iurino D.A., Menconero S., Bona F., Cirilli O., Kimambo J.S., Masao F.T., Barili A., Ambrose S.H. & Manzi G.	29
Old World Early Pleistocene Canidae record: a biogeographic perspective <i>by Bartolini Lucenti S.</i>	30
DISSCO RI: towards a European infrastructure for scientific collections <i>by Bellucci L.</i> , Bartolozzi L., Biaggini M., Cecchi L., Di Vincenzo F., Innocenti G., Manca R., Mancinelli M.L., Moggi Cecchi V., Porena M., Presutti V., Rossi P.F., Rossi De Gasperis S., Veninata C. & Benvenuti M.	31
Middle Triassic: the onset of modern terrestrial tetrapod faunas <i>by Bindellini G.</i> , Benton M.J., Petti F.M., Bernardi M., Rubidge B., Hancox J. & Romano M.	32
Neogene cold-seep microbial carbonates (Crotone basin - South Italy) <i>by Borrelli M.</i> , Perri E., Heimhofer U., Santagati P. & Le Pera E.	33
Eocene stilt walkers: unique rotatory growth strategy of the scleractinian coral <i>Stylocoenia</i> <i>by Bosellini F.R.</i> & Stolarski J.	34

Cuticles: a world waiting to be discovered	35
<i>by Branz R. & Kustatscher E.</i>	
The Rhamphosidae (Teleostei, Syngnathiformes) from the Eocene of Bolca	36
<i>by Calzoni P., Amalfitano J., Marramà G., Carnevale G. & Giusberti L.</i>	
The dawn of moorish idols (Teleostei, Zanclidae) in the Eocene of Bolca	37
<i>by Carnevale G. & Tyler J.C.</i>	
Are the Mississippian mud-dominated carbonate buildups of Derbyshire “mud mounds”?	38
<i>by Carniti A.P., Della Porta G. & Angiolini L.</i>	
Risposta delle associazioni a foraminiferi alle oscillazioni climatiche durante il Gelasiano-Calabriano: dati ottenuti dalla successione di Capo Rossello (Sicilia)	39
<i>by Caruso A., Cosentino C., Scopelliti G. & Herbert T.</i>	
Tempo e modo nell’evoluzione della nicchia climatica in Dinosauria (Sauropsida: Archosauria)	40
<i>by Chiarenza A.A., Farnsworth A.J., Jones L.A., Cantalapiedra J.L., Gamboa S., Galvan S., Sotelo G. & Varela S.</i>	
Cenozoic carbonate facies of Asia: the hidden potential of the shallow water record	41
<i>by Coletti G., Commissario L., Mariani L., Bosio G., Desbiolles F., Soldi M. & Bialik O.M.</i>	
Mollusc shells as a tool to reconstruct the palaeoenvironment and the palaeoclimate in the HAS1 settlement (Holocene, Oman)	42
<i>by Crippa G., Chiari A., Cremaschi M., Dapiaggi M., Lischi S. & Leng M.J.</i>	
Double-phased biomineralization processes in marine invertebrates: the example of cirratulids (Polychaeta, Annelida) from southern Peru	43
<i>by D’Amico F., Guido A., Kočí T., DeVries T.J., Collareta A., Bosio G. & Sanfilippo R.</i>	
Seasonal growth patterns of pectinids from the warm Mediterranean Pliocene	44
<i>by Danise S., Giachetti G., Baneschi I., Casalini M., Miniati F., Dominici S. & Boschi C.</i>	
Morphometric analyses of the Pliocene oysters from Buttiglieri d’Asti (Asti, NW Italy)	45
<i>by De Bortoli L.</i>	
New alloodaposuchid remains from the Late Cretaceous of Catalonia (Spain): insights into dermal and postcranial skeletal anatomy	46
<i>by Della Giustina F. & Vila B.</i>	
DeepBryo: a web app for bryozoan AI-assisted morphometric characterization	47
<i>by Di Martino E. & Porto A.</i>	

The silicified molluscs of Pelsa-Vazzoler (eastern Dolomites, Italy) and the late Middle Triassic diversification of marine life	48
<i>by Dominici S.</i> , Danise S. & Tintori A.	
Assessing relevance and vulnerability of palaeontological sites: an analytical procedure	49
<i>by Faggi A.</i> , Bartolini Lucenti S. & Rook L.	
60 years of the Italian Paleontological Society through its journal	50
<i>by Ferretti A.</i> & Balini M.	
Reassessment of a putative peltopleurid fish from the Sinemurian (Lower Jurassic) of Osteno (Como, Italy): preliminary results	51
<i>by Franceschi F.</i> , Marramà G. & Carnevale G.	
The analysis of the frontal sinus in the genus <i>Eucyon</i> : evolutionary and ecological insights	52
<i>by Frosali S.</i> , Bartolini-Lucenti S., Madurell-Malapeira J., Costeur L. & Rook L.	
Microbial biomineralizations in tidally-influenced subterranean settings (Zinzulùsa Cave, Castro, Italy)	53
<i>by Guido A.</i> , Alifano P., Talà A., Calcagnile M. & Belmonte G.	
Bethlehem suids reconsidered: chronological and paleobiogeographic implications	54
<i>by Iannucci A.</i> , Souron A. & Rabinovich R.	
The first Kungurian fossil forest reveals wetland conifers to thrive in peri-Tethyan Pangaea	55
<i>by Kustatscher E.</i> , Trümper S., Rößler R., Morelli C., Krainer K., Karbacher S., Vogel B., Antonelli M. & Sacco E.	
<i>Plioviverrops faventinus</i> , a small Messinian hyaenid from Italy: a revision	56
<i>by Loddi C.</i> , Bartolini Lucenti S., Madurell-Malapeira J. & Rook L.	
<i>Pachycrocuta</i> , an active hunter or a social scavenger? Insights from postcranial morphology	57
<i>by Madurell-Malapeira J.</i> , Bartolini Lucenti S., Prat-Vericat M., Costeur L. & Rook L.	
Coccolithophore variations in the subantarctic South Pacific along late Quaternary glacial-interglacial cycles	58
<i>by Malinverno E.</i> , Saavedra-Pellitero M., Colucci M. & IODP-383 Scientists	
Double-armoured herrings (Clupeomorpha: Ellimmichthyiformes) from the Lower Cretaceous of Pietraroja (Southern Italy)	59
<i>by Marramà G.</i> & Carnevale G.	
New paleontological results on the early Middle Pleistocene of Notarchirico (Venosa, Southern Italy)	60
<i>by Mecozzi B.</i> , Iannucci A., Sardella R., Carpentieri M., Pineda A., Rabinovich R. & Moncel M.H.	

It's time to eat: great white shark (<i>Carcharodon carcharias</i>) bite marks on balaenid (right whale) remains from the Pliocene of Tuscany (Central Italy)	61
<i>by Merella M.</i> , Collareta A., Casati S., Di Cencio A., Pieri A. & Bianucci G.	
Bivalve response to Plio-Pleistocene climate change in the Mediterranean Sea	62
<i>by Mondanaro A.</i> , Dominici S. & Danise S.	
Endocranial pneumatization of <i>Ursus spelaeus</i> and implications in the bite force	63
<i>by Moscarella A.</i> , Iurino D.A., Conti J., Bona F., Di Giancamillo M. & Sardella R.	
Synchrotron tomography provides novel insights into the dental anatomy of <i>Acynodon adriaticus</i> (Eusuchia, Hylaeochampsidae)	64
<i>by Muscioni M.</i> , Dreossi D. & Fanti F.	
Cambrian bipartite animals reveal a lost chapter in chordate history	65
<i>by Mussini G.</i> , Smith M.P., Rahman I.A., Murdock D.J.E., Vinther J., Harper D.A.T. & Dunn F.S.	
First Mediterranean record of the delphinidan genus <i>Kentriodon</i> from a new lower Miocene odontocete locality at Feltre (northeastern Italy)	66
<i>by Nobile F.</i> , Collareta A., Perenzin V., Fornaciari E., Giusberti L. & Bianucci G.	
Deconstructing the classical Late Miocene faunal assemblages of the Tusco-Sardinian paleobioprovince	67
<i>by Pandolfi L.</i> & Rook L.	
Middle Pleistocene to Holocene ostracod assemblages from the Corinth Gulf	68
<i>by Parisi R.</i> & Mazzini I.	
The outstanding <i>Dama</i> -like deer sample from Vallparadís Estació (Iberian Peninsula, Early Pleistocene)	69
<i>by Parparousi E.M.</i> , Madurell-Malapeira J., Sorbelli L., Breda M. & Cherin M.	
Microbially mediated Ca-carbonate precipitation in tufa fluvial system: an innovative approach to develop a new generation of sustainable bio-inspired materials	70
<i>by Perri E.</i> , Borrelli M. & Ponte M.	
<i>In-vivo</i> 3D reconstruction and first volumetric body mass estimate of the oldest Karoo pareiasaur <i>Bradyosaurus baini</i>	71
<i>by Romano M.</i> , Van den Brandt M.J., Day M.O., Manucci F., Viglietti P.A. & Angielczyk K.D.	
Impossible fossils: can soft tissues fossilize in volcaniclastic settings?	72
<i>by Rossi V.</i> , Iurino D.A., Terranova E., Fornasiero M., Bernardi M., Sardella R. & Kustatscher E.	

Biogenic content of surface cores from deep-shelf bottoms close to the coralligenous off Marzamemi (SE Sicily, Ionian Sea): preliminary data	73
by Rosso A. , Sanfilippo R., Donato G., Sciuto F., D'Alpa F., Caccamo D., Minniti G., Bracchi V.A., Savini A., Malinverno E., Guido A., Bazzicalupo P., Negri M.P., Cipriani M., Leonardi R., Viola A. & Basso D.	
MAE: Museo Archeologico Ernico - A brand-new museum in Anagni is born	74
by Sardella R. , Conti J., Germani M., Iannucci A., Iurino D.A., Mecozzi B., Messina M.Y., Moscarella A., Pioggia M., Strani F. & Vona S.	
<i>Leptobos</i> and <i>Bison</i> limb proportions and body mass changes in response to environmental transitions during the Pleistocene	75
by Sorbelli L. , Madurell-Malapeira J., Brugal J.P., Kostopoulos D.S. & Cherin M.	
Rediscovering the fossil collection from the late Early Pleistocene site of Pirro Nord (Foggia, Southern Italy) at the Bari University	76
by Stefanelli D. , Mecozzi B., Breda M., Girone A., La Perna R., Maiorano P., Marino M. & Sardella R.	
Combining palaeoecological and palaeoneurological analyses to disentangle cervid evolutionary history in relation to climate changes: the case study of the Early Pleistocene deer from Pirro Nord (Italian Peninsula)	77
by Strani F. , Di Folco F., Iurino D.A., Cherin M., Pushkina D., Rook L., Sardella R., Azanza B. & DeMiguel D.	
Collezioni storiche: problematiche e strategie per la valorizzazione e comunicazione nel caso della Collezione Georg Gasser (1857-1931)	78
by Tomelleri I. , Wagensommer A. & Kustatscher E.	
Reconstruction of depositional environments in the Kungurian Megacaldera of the Athesian volcanic group: a multi-proxy approach	79
by Vallè F. , Nowak H., Kustatscher E., Roghi G., Morelli C., Krainer K., Preto N. & Hartkopf-Fröder C.	
Silicified shells as archives of geochemical proxies: the hidden potential of a diagenetic process	80
by Viaretti M. , Jurikova H. & Angiolini L.	
<hr/> POSTERS	<hr/> 81
Multidisciplinary analysis of the sedimentary succession at San Giuliano Lake (Matera, Southern Italy): sedimentological, biostratigraphic and palaeoecological reconstruction	82
by Amore F.O. , Argenio C., Faranda C., Ferraro L., Gliozi E., Magri D., Michelangeli F., Relvini A., Russo B., Siciliano J., Vallefuoco M., Meo A. & Senatore M.R.	
Theropod tracks from the Lama Badessa ichnosite (middle-upper Cenomanian, Apulia, Southern Italy)	83
by Antonelli M. , Petti F.M., De Sario F., Sacco E., Petruzzelli M., Sabato L., Spalluto L. & Tropeano M.	

Disparity vs diversity within the <i>Hemisyntrachelus</i> lineage (Cetacea, Delphinidae): a review of the Italian Pliocene record	84
<i>by Bianucci G.</i> , Geisler J.H., Collareta A., Sfravara L., Merella M., Carnevale G., Damarco P. & Ferretti M.P.	
Impact of plastic pollution on benthic foraminifera	85
<i>by Birarda G., Buosi C.</i> , Caridi F., Casu M.A., De Giudici G., Di Bella L., Medas D., Meneghini C., Pierdomenico M., Sabbatini A., Surowka A. & Vaccari L.	
Eccentricity signal in the time-series of nannofossil species at mid-latitude across the mid-Pleistocene transition (ODP site 1209, northwestern Pacific Ocean)	86
<i>by Bordiga M.</i> , Lupi C., Sacchi R., Ferretti P., Crowhurst S.J. & Cobianchi M.	
Paleoenvironmental considerations on the Val Seriana Valley (north-western Italy) during the Upper Pleistocene. The small-mammal fossils from Büs di Tri Fradei Cave (Oltre il Colle, BG) as palaeoclimatic proxy data	87
<i>by Carnevali V. & Bona F.</i>	
Brachiopods and conodonts from the upper Mississippian Tonka Formation of Carlin Canyon, Nevada	88
<i>by Carniti A.P.</i> , Anderson C.M. & Angiolini L.	
Odontometric characteristics of the European Miocene primates	89
<i>by Cieszynska W.K.</i>	
Evidence of not-genetically controlled biomineralizations in coralligenous build-ups (Marzamemi, Sicily, Italy)	90
<i>by Cipriani M., Apollaro C., Basso D., Bazzicalupo P., Bertolino M., Bracchi V.A., Bruno F., Costa G., Dominici R., Gallo A., Muzzupappa M., Rosso A., Sanfilippo R., Sciuto F., Vespasiano G. & Guido A.</i>	
Mesozoic and Cenozoic carbonate platform bioevents from the CARG Geological Sheet n. 377 “Trasacco”	91
<i>by Consorti L., Fabbri S.</i> , Cipriani A. & Pampaloni M.L.	
Agents of climate change: a paleontological point of view	92
<i>by D’Amico F.</i> , Calidonna C.R., Apollaro C., Vespasiano G. & Guido A.	
I resti di coccodrilli tomistomini della Pietra Leccese (Miocene, Italia meridionale)	93
<i>by Delfino M.</i> , Amoruso N., Macaluso L. & Rook L.	
The Cretaceous flora of Salento: preliminary data on the assemblage from Surbo (Lecce, Italy)	94
<i>by De Pirro D.</i> , Bartiromo A., Cherin M., Lionello P. & Spina A.	
Live and dead bryozoans in the coralligenous build-ups of Marzamemi (Ionian Sea)	95
<i>by Donato G.</i> , Sanfilippo R., D’Alpa F., Sciuto F., Bracchi V.A., Basso D. & Rosso A.	

New insights on Middle Triassic invertebrate ichnoassociation from the Monti Pisani megaichnosite (northern Apennines)	96
<i>by Granata V. & Collareta A.</i>	
Microfacies and biosedimentary processes of Late Pleistocene (MIS 5.1) deposits (Capo Colonna, Southern Italy)	97
<i>by Guerrieri S., Santagati P., Borrelli M. & Perri E.</i>	
A flat-snouted sperm whale from the Lower Miocene of the southeastern Pacific (east Pisco Basin, Peru) supports affinities with the southwestern Atlantic cetacean fauna	98
<i>by Lambert O., Collareta A., Benites-Palomino A., Merella M., de Muizon C., Bennion R., Urbina M. & Bianucci G.</i>	
New fossil remains and histological study of <i>Diplocynodon ratelii</i> (Crocodylia: Diplocynodontidae) from the Middle Miocene site of Els Casots (Catalonia)	99
<i>by Logrosán Soriano P., Luján A.H., Nacarino-Meneses C., Villa A., Abella J., Madurell-Malapeira J. & Casanova-Vilar I.</i>	
Analisi paleoecologica dell'associazione a <i>Venus-Ostrea</i> del Torrente Arda (Pleistocene Inferiore)	100
<i>by Lombardi M. & Crippa G.</i>	
Revising the early European record of the dirk-toothed cat <i>Megantereon</i>	101
<i>by Madurell-Malapeira J., Bartolini-Lucenti S., Costeur L. & Rook L.</i>	
Paleoclimatic and paleoceanographic changes and their impact on planktic foraminifera in the Tyrrhenian Sea during the last 32 ka	102
<i>by Martinelli P., Gennari R., Lirer F. & Cascella A.</i>	
Technological innovation for the remote 3D study of marine bioconstructions: a project proposal	103
<i>by Maruca G., Bruno F., Lagudi A., Barbieri L., Gallo A., Cipriani M. & Guido A.</i>	
The Pleistocene malacological collection of Catallarga (Sicily) in the Museum of Paleontology (University of Catania)	104
<i>by Minniti G., Reitano A., Di Geronimo I., Rosso A. & Sanfilippo R.</i>	
Digital investigation of lamniform shark vertebrae from the Scaglia Cinerea – Bisciaro boundary (Miocene, Umbria-Marche basin)	105
<i>by Moscarella A., Fabbi S., Bindellini G., Carnevale G., Citton P., Consorti L., D'Ambrogi C., Garzarella A., Muraro C., Pampaloni M.L., Prinzi E.P., Radeff G., Romagnoli G., Romano M., Spanò F. & Cipriani A.</i>	
Preliminary study of two cave bear cubs from the Late Pleistocene vertebrate assemblage of the “Tecchia” of Equi cave (Equi Terme, Massa-Carrara province, Italy)	106
<i>by Mulè F., Collareta A. & Farina S.</i>	

A leathery meal: megatoothed shark bite mark on a <i>Psephophorus</i> carapace from the Pietra Leccese formation (Miocene, Southern Italy) and its palaeoecological implications	107
<i>by Peri E.</i> , Collareta A., Granata V. & Bianucci G.	
The Wolferode Quarry: an amazing middle to late Permian tracksite in Central Germany	108
<i>by Pieri A.</i> , Falk D., Buchwitz M., Schneider J.W. & Ehling B.C.	
Moleto: a new latest middle-earliest Late Pleistocene vertebrate site in Northern Italy	109
<i>by Piñero P.</i> , López-García J.M., Carnevale G., Furió M., Giuntelli P., Luzi E., Macaluso L., Marramà G., Pavia G., Pavia M., Pezzetti C., Rocca M., Villa A. & Delfino M.	
Nuove evidenze di depositi plio-pleistocenici a conchiglie e vegetali fossili nell'alveo del Torrente Cervo: rassegna preliminare	110
<i>by Pizzitola V.O.</i> & Martinetto E.	
Reimagining <i>Velociraptor</i> 100 years after its discovery	111
<i>by Romano M.</i> , Bindellini G., Moscarella A., Manucci F. & O'Connor J.K.	
The Achille Tellini historical collection (Museo Universitario di Scienze della Terra, Sapienza Università di Roma)	112
<i>by Sacco E.</i> , Pignatti J.S. & Sardella R.	
Biodiversity and distribution pattern of serpulids from Mediterranean Holocene coralligenous build-ups	113
<i>by Sanfilippo R.</i> , Donato G., Basso D., Bracchi V.A. & Rosso A.	
MIS 5.5 SST estimation from a macrofossil assemblage: a multi-proxy reconstruction	114
<i>by Santagati P.</i> , Perri E., Borrelli M. & Guerrieri S.	
Collezioni museali e valorizzazione del patrimonio paleontologico. Mappare per costruire reti: il caso studio della Val d'Alpone	115
<i>by Tomelleri I.</i> & Zorzin R.	
A dendrolite/thrombolite carbonate build-up from the Messinian terminal carbonate complex of the Salento Peninsula (Southern Italy)	116
<i>by Vescogni A.</i> & Guido A.	
Some considerations about MIS3 marmots (<i>Marmota marmota</i>) of north-western Italy with particular regards to those from Caverna Generosa (Centrovalle Intelvi, Como)	117
<i>by Vincenzi M.</i> , Bona F. & Nappi A.	
Re-animation of a fossil: bringing back to life the specimens from MUSTE (Museo di Scienze della Terra) of University of Bari with digital technologies	118
<i>by Zazzera A.</i> , Marsico A., La Perna R., Marino M., Venisti N., De Prezzo G., Bruno Stamerra F., Spisso N., Spada I. & Girone A.	
A new proboscidean tracksite in the Pleistocene of south-western Sardinia (Italy)	119
<i>by Zoboli D.</i> , Masala A. & Pillola G.L.	