

# Acknowledgements

This research was supported by the Università del Salento, by the “Università Italo Francese”/“Université Franco Italienne”(UIF/ UFI) under the Mobility grant of VINCI 2008 (Funding for joint PhD between Italy and France), by the “Centre National de la Recherche Scientifique” and by the “Gruppo Nazionale per la Fisica Matematica” of Italian Istituto Nazionale di Alta Matematica.

I would like to express my thanks to the people who have helped me during the time it took me to write this Thesis.

First and foremost, my gratitude goes to Michel Destrade and to Giuseppe Saccomandi, who I really thank for having been my supervisors, and who complemented each other wonderfully well.

I am also grateful to: Martine Ben Amar (Paris), Alain Goriely (Tucson-Oxford), Corrado Maurini (Paris), Giorgio Metafuno (Lecce), Gaetano Napoli (Lecce), Ray W. Ogden (Glasgow), Diego Pallara (Lecce), Kumbakonam R. Rajagopal (College Station), Ivonne Sgura (Lecce), Raffaele Vitolo (Lecce).

Special Gratitude goes to the Institut Jean Le Rond d’Alembert and all the people of the Laboratory for having welcomed me during my second year of PhD and my long visit to Paris.

Finally, I do not forget all the support I received from my friends in Lecce and in Paris and from my family, many thanks.